

Gender Development and the Human Brain

Melissa Hines

Department of Social and Developmental Psychology, University of Cambridge,
Cambridge, CB2 3RQ, United Kingdom; email: mh504@cam.ac.uk

Annu. Rev. Neurosci. 2011. 34:69–88

First published online as a Review in Advance on
March 25, 2011

The *Annual Review of Neuroscience* is online at
neuro.annualreviews.org

This article's doi:
10.1146/annurev-neuro-061010-113654

Copyright © 2011 by Annual Reviews.
All rights reserved

0147-006X/11/0721-0069\$20.00

Keywords

sex difference, testosterone, behavior, sex differentiation, androgen,
play

Abstract

Convincing evidence indicates that prenatal exposure to the gonadal hormone, testosterone, influences the development of children's sex-typical toy and activity interests. In addition, growing evidence shows that testosterone exposure contributes similarly to the development of other human behaviors that show sex differences, including sexual orientation, core gender identity, and some, though not all, sex-related cognitive and personality characteristics. In addition to these prenatal hormonal influences, early infancy and puberty may provide additional critical periods when hormones influence human neurobehavioral organization. Sex-linked genes could also contribute to human gender development, and most sex-related characteristics are influenced by socialization and other aspects of postnatal experience, as well. Neural mechanisms underlying the influences of gonadal hormones on human behavior are beginning to be identified. Although the neural mechanisms underlying experiential influences remain largely uninvestigated, they could involve the same neural circuitry as that affected by hormones.

Contents

INTRODUCTION	70
GENERAL PRINCIPLES OF SEXUAL DIFFERENTIATION	70
ANIMAL MODELS OF HORMONE EFFECTS	71
HUMAN RESEARCH	72
Why Study Children's Play?	72
Is the Behavioral Alteration Caused by Hormones Acting on the Developing Brain?	73
Rethinking Children's Preferences for Sex-Typed Toys	73
Early Hormone Influences on Sexual Orientation and Core Gender Identity	74
Early Hormone Influences on Personality and Cognition	75
Socialization and Sex Differences in Cognitive Performance	76
Sex-Related Psychiatric Disorders	76
Sex Differences in Brain Structure and Function	77
Effects of Experience on the Brain	78
Other Potential Critical Periods for Hormone Influences on Gender Development	79
CONCLUDING REMARKS	80

INTRODUCTION

Males and females differ both behaviorally and neurally. Indeed, the existence of behavioral sex differences implies the existence of neural sex differences, given that behavior depends on the nervous system. Contemporary research shows that gendered behavior results from a complex interplay of genes, gonadal hormones, socialization, and cognitive development related to gender identification. This article focuses on the role of gonadal hormones, particularly testosterone, during early development. This focus has been chosen because extensive experimental research in nonhuman mammals shows that testosterone exerts powerful

influences on both gender-related behavior and the developing brain and because recent research provides convincing evidence that testosterone exerts similar influences on human development. The article critically reviews the evidence of prenatal hormonal influences on human neurobehavioral sexual differentiation and contextualizes these hormonal effects with genetic, social, and cognitive influences on gender development. It also critically reviews the evidence regarding possible neural changes underlying hormonal influences on human behavior and suggests that neural systems similar to those influenced by hormones underlie other types of influences on gendered behavior.

GENERAL PRINCIPLES OF SEXUAL DIFFERENTIATION

Gender development begins at conception with the union of two X chromosomes (genetic female) or an X and a Y chromosome (genetic male). The main role of these sex chromosomes in human sexual differentiation is to determine whether the gonads become testes or ovaries (Arnold 2009). Genetic information on the Y chromosome leads to testicular differentiation (Wilson et al. 1981), whereas without the Y chromosome, ovaries develop instead of testes. The testes begin to produce testosterone prenatally, and the ovaries do not (Wilson et al. 1981). Consequently, male and female fetuses differ in the amount of testosterone to which they are exposed. This sex difference appears to be maximal between about weeks 8 and 24 of human gestation, with testosterone in males tapering off before birth (Carson et al. 1982, Reyes et al. 1973). In nonhuman mammals at comparable stages of early development, testosterone and hormones produced from testosterone influence neural survival, neuroanatomical connectivity, and neurochemical specification, producing sex differences in brain structure and function (McCarthy et al. 2009). These effects of testosterone on neural development provide powerful mechanisms for influencing behavior across the life span.

Gonadal hormones:

gonads' products, including androgens, produced mainly by the testes, and estrogens and progesterone, produced mainly by the ovaries

Testosterone: the major androgenic hormone produced by the testes

ANIMAL MODELS OF HORMONE EFFECTS

The influences of early testosterone exposure on neurobehavioral development were first documented in a landmark study by Phoenix et al. (1959). They showed that administering testosterone to pregnant guinea pigs produced female offspring who showed increased capacity for male-typical sexual behavior and decreased capacity for female-typical sexual behavior in adulthood (Phoenix et al. 1959). Phoenix et al. contrasted these early, and permanent, effects of hormones, which they called organizational because they were thought to reflect changes in the organization of neural systems, with the later, and transient, effects of hormones after puberty, which they called activational because they were thought to reflect transient activation of the previously organized systems. This organizational/activational distinction has stood up well in the subsequent 50 some years (Arnold 2009), and thousands of studies on numerous species, including not only guinea pigs, but also rats, mice, hamsters, gerbils, ferrets, dogs, sheep, and marmoset and rhesus monkeys, have documented the early organizing effects of testosterone on a wide variety of behaviors that show sex differences (Hines 2004, 2009; McCarthy et al. 2009). For instance, the female offspring of rhesus macaques treated with testosterone during pregnancy show increased male-typical, and reduced female-typical, sexual behavior in adulthood, and increased male-typical, rough-and-tumble play as juvenile animals.

The organizing influences of testosterone on behavioral development were originally thought to reflect subtle neural changes (Phoenix et al. 1959). Subsequent research, however, has shown that early hormone manipulations produce dramatic changes in the structure of neural regions with the relevant hormone receptors. The first dramatic neural sex difference described in the rodent brain was the sexually dimorphic nucleus of the preoptic area (SDN-POA). This region of the anterior hypothalamic/preoptic area (AH/POA) is several

fold larger in the adult male rat than in the adult female rat, and its volume can be altered by manipulating testosterone during early development (Gorski et al. 1978, 1980). Administering testosterone to developing female animals increases the volume of the SDN-POA, and removing testosterone from developing males reduces its volume (Dohler et al. 1984, Jacobson et al. 1981). Other neural regions in addition to the SDN-POA show sex differences, and in these regions too, the size of the sex difference is influenced by the early hormone environment. For instance, a second region of the preoptic area, the anteroventral paraventricular nucleus, is larger and contains more neurons in female rats than in males, and these characteristics are reduced by early testosterone treatment (Ito et al. 1986, Sumida et al. 1993). Similar neural sex differences have been reported in other rodent species, including gerbils, hamsters, mice, and guinea pigs, as well as in ferrets, sheep, and rhesus monkeys, and studies investigating early hormone influences have found similar results to those seen in rats, in other rodent species, and in ferrets (Bleier et al. 1982, Byne 1998, Hines et al. 1987, Roselli et al. 2004, Simerly et al. 1997, Tobet et al. 1986, Ulibarri & Yahr 1988).

Some general principles can be derived from the extensive experimental work in nonhuman mammals, and these principles have informed hypotheses regarding possible hormonal influences on human brain and behavior (see Hines 2009, for a review). First, during early development, estrogens generally do not promote female-typical development. Instead, female-typical development occurs in the absence of testicular hormones. Thus, exposure to high levels of estrogen is not expected to femininize neurobehavioral development. Second, the effects of testosterone on development are graded and linear; the more hormone the animal is exposed to, the more male-typical its behavior and brain structure become. An implication of this principle is that gonadal hormones can contribute to individual differences within each sex, as well as to differences between the sexes. Third, neurobehavioral sexual differentiation is a multidimensional process. The many

Rough-and-tumble play: juvenile behavior characterized by overall body contact or playful aggression; more common in males than in females

SDN-POA: sexually dimorphic nucleus of the preoptic area

Critical period:

programmed stage of development at which an influence is most likely (or only likely) to occur. Sometimes called sensitive period

Androgenic**(anti-androgenic)**

progesterins: synthetic hormones that mimic progesterone, but which can also mimic androgens (androgenic) or impair androgen action (antiandrogenic)

Congenital adrenal**hyperplasia (CAH):**

genetic disorder causing increased adrenal androgen production, beginning prenatally

behaviors and neural systems that differ for males and females can be influenced by hormones during slightly different critical periods, or can be sensitive to different doses of hormone, or to different metabolites of testosterone, or can involve different downstream mechanisms such as cofactors. Implications of this principle include an expectation that the many human behaviors and brain structures that differ by sex may not relate in a uniform way to one another and that individuals can develop complicated patterns of sex-typed behavior, being masculine in some respects and feminine in others. Fourth, the effects of hormones can differ somewhat from one species to another. For instance, behaviors that are influenced in one species may not be influenced in all others. Similarly, brain regions that differ for males and females in one species may not show a sex difference in another. Thus the specific effects of gonadal steroids seen on the brain and behavior of nonhuman mammals cannot be automatically generalized to humans, as well. Instead, hypothesized neural and behavioral influences of testosterone during early development must be evaluated directly in humans. Fifth, the behaviors and neural features that are influenced by gonadal hormones are those that show sex differences, meaning that they differ on average for males and females. Therefore, the characteristics that are likely to be influenced in humans are also those that show sex differences.

HUMAN RESEARCH

Ethical considerations generally preclude experimental manipulations of gonadal hormones in humans during early development. However, information from genetic syndromes that produce fetal hormone abnormality, as well as from situations in which pregnant women have been prescribed hormones, and studies relating normal variability in hormones early in life to normal variability in subsequent behavior all suggest that hormones contribute to human gender development. The most convincing evidence of these influences has come from studies of childhood play.

Why Study Children's Play?

Girls and boys differ in their toy, playmate, and activity preferences (Hines, 2010a). For example, boys tend to prefer toy vehicles, whereas girls tend to prefer dolls. Girls and boys also generally prefer playmates of their own sex, and boys spend more time in rough-and-tumble play than girls do. Children's sex-typed play behavior is the aspect of human gender development that has been studied most extensively in relation to the early hormone environment. This focus on childhood play reflects several considerations. First, children spend most of their time playing, and play is thought to be essential for healthy cognitive and emotional development (Ginsburg et al. 2007, Piaget 1970, Vygotsky 1976). Second, children's sex-typed play behavior can be assessed readily and reliably. Third, large sex differences exist in children's play, larger than those in cognitive abilities or personality characteristics (Hines 2010b), providing scope for detecting hormonal influences. Fourth, sex differences in children's play are evident early in life and relate to other behaviors that show sex differences, including sexual orientation and gender identification (Bailey & Zucker 1995, Green 1985, Hines et al. 2004). Fifth, play can be assessed during a period of hormonal quiescence, allowing examination of the early and permanent organizational influences of hormones on brain development, prior to the addition of the transient, activational influences of hormones that occur after puberty.

Several types of studies provide convergent evidence that testosterone concentrations prenatally influence children's subsequent sex-typed toy, playmate, and activity preferences. Studies of girls exposed to unusually high levels of testosterone and other androgens before birth, because they have the genetic disorder known as classic congenital adrenal hyperplasia (CAH), consistently find that these girls show increased male-typical play and reduced female-typical play (Berenbaum & Hines 1992, Dittmann et al. 1990, Ehrhardt et al. 1968, Ehrhardt & Baker 1974, Hall et al.

2004, Nordenstrom et al. 2002, Pasterski et al. 2005). Similarly, children whose mothers took androgenic progestins during pregnancy have shown increased male-typical toy and activity preferences, whereas the opposite occurs in children whose mothers took antiandrogenic progestins (Ehrhardt et al. 1977, Ehrhardt & Money 1967).

Is the Behavioral Alteration Caused by Hormones Acting on the Developing Brain?

The external genitalia, as well as the brain, contain androgen receptors, and girls with CAH, as well as those whose mothers took androgenic progestins, are typically born with varying degrees of genital virilization (enlarged clitoris, fused labia). Those skeptical of gonadal hormone influences on human neurobehavioral development suggest that the abnormal genital appearance, rather than the neural influences of androgens, could cause behavioral masculinization (Fausto-Sterling 1992, Jordan-Young 2010). Specifically, they suggest that parents may treat their daughters differently because of the girls' external virilization at birth and that this difference in parental treatment could alter sex-typed behavior. In addition, they suggest that virilized genitalia could reduce self-identification as female, which could in turn cause increased male-typical behavior.

Some evidence suggests that parents can influence the development of children's gender-typical behavior. For instance, parents generally encourage sex-typical play (Fagot 1978, Langlois & Downs 1980, Pasterski et al. 2005), and the amount of such encouragement has been found to correlate with the amount of sex-typed toy play, at least among typically developing children (Pasterski et al. 2005). However, parents have been found to offer more, rather than less, encouragement of sex-typical play to their daughters with CAH than to their daughters who do not have the disorder (Pasterski et al. 2005), suggesting that parental encouragement is not responsible for cross-gendered toy choices in girls with CAH.

Similarly, although gender identification plays a role in children's acquisition of gender-related behavior, at least in typically developing children (Hines 2010a, Ruble et al. 2006), it is unlikely that the male-typical behavior in girls with CAH results solely from altered gender identity based on genital virilization at birth. Evidence arguing against this explanation comes from studies relating normal variability in prenatal testosterone exposure to normal variability in subsequent behavior. Testosterone concentrations in maternal blood samples taken during pregnancy or in amniotic fluid from normally developing fetuses relate positively to male-typical childhood behavior (Auyeung et al. 2009b, Hines et al. 2002). Because the children in these studies have normal appearing genitalia, it is unlikely that differential parental socialization or changes in gender identification based on genital appearance account for the observed relationships between prenatal testosterone and postnatal behavior.

Researchers have also looked at species in which children's toys are novel objects and therefore not subject to the socialization histories or processes of gender identification thought to explain sex-typed toy preferences in children. Two studies of nonhuman primates have reported sex-typed toy preferences similar to those seen in children. Male rhesus monkeys (Alexander & Hines 2002) have been found to spend more time than females contacting toys that are typically preferred by boys (e.g., a car) and less time contacting toys that are typically preferred by girls (e.g., a doll) (**Figure 1**). Similarly, male rhesus monkeys have been found to prefer toys normally preferred by boys (wheeled toys) to plush toys (Hassett et al. 2008). These findings show that sex-typed toy preferences can arise independent of the social and cognitive processes involved in gender development.

Rethinking Children's Preferences for Sex-Typed Toys

Children's sex-typical toy preferences have been widely assumed to result from socialization and other postnatal factors and to

Androgens:

substances, including testosterone, that promote masculinization. Produced by the testes, adrenal glands, and ovaries, with the testes the largest source

Virilized genitalia:

masculinized genitalia, typically involving an enlarged clitoris and partially fused labia



Figure 1

Examples of a male and a female vervet monkey contacting human children's sex-typed toys. The female animal (*left*) appears to be inspecting the doll in a manner similar to that in which vervet monkeys inspect infant vervets. The male animal (*right*) appears to be moving the car along the ground as a child might do. Reproduced by permission from Alexander & Hines (2002).

provide rehearsals for adult sex-typed social roles. Evidence of inborn influences has led researchers to reevaluate this perspective and to investigate the object features that make certain toys more or less interesting to brains exposed prenatally to different amounts of testosterone. Although boys' toys and girls' toys differ in shape and color (boys' toys tend to be angular and blue, whereas girls' toys tend to be rounded and pink), sex differences in toy preferences are present in very young infants (Alexander et al. 2009b, Campbell et al. 2000, Jadva et al. 2010, Serbin et al. 2001), before sex differences in color or shape preferences are seen (Jadva et al. 2010), suggesting that the object preferences do not result from the color or shape preferences. Another possibility is that boys like toys that can be moved in space,

and prenatal androgen exposure may increase interest in watching things move in space (Alexander 2003, Alexander & Hines 2002, Hines 2004), perhaps by altering development of the visual system (Alexander 2003).

Early Hormone Influences on Sexual Orientation and Core Gender Identity

Adult behaviors that show sex differences, including sexual orientation and core gender identity, also appear to be influenced by prenatal testosterone exposure. Women with CAH not only recall more male-typical childhood behavior, but also show reduced heterosexual orientation as adults, and these two outcomes correlate (Hines et al. 2004; see Meyer-Bahlburg et al. 2008 for a review of additional

studies of CAH and sexual orientation). Normal variability in testosterone prenatally, e.g., from maternal blood or amniotic fluid, has not yet been related to sexual orientation, but a characteristic that is thought to provide an indirect measure of prenatal testosterone exposure, the ratio of the second to the fourth digit of the hand (2D:4D), which is greater in females than in males, has been linked. A study of more than 200,000 individuals, who measured their own 2D:4D and reported their sexual orientation online, found that 2D:4D related as predicted to sexual orientation in males, but not in females (Collaer et al. 2007). A meta-analysis that did not include this large study reached a somewhat different conclusion, however, finding that 2D:4D related as predicted to sexual orientation in females, but not in males (Grimbos et al. 2010). Finger ratios are probably a weak correlate of prenatal testosterone exposure, perhaps explaining the somewhat inconsistent results.

Women with CAH not only show reduced heterosexual interest, but also show diminished identification with the female gender, and this too correlates with their recalled childhood sex-typical behavior (Hines et al. 2004; see Hines 2010a for a review of studies of gender identity in females with CAH). About 3% of women with CAH express a desire to live as men in adulthood, despite having been reared as girls, in contrast with ~0.005% of all women (Dessens et al. 2005). Although 3% may seem small, it indicates that women with CAH are ~600 times more likely than women in general to experience severe gender dysphoria. Girls with other disorders involving exposure to unusually high levels of androgens prenatally also show increased gender dysphoria (Slijper et al. 1998). Additionally, even when not gender dysphoric, or wishing to change sex, girls and women with CAH show somewhat reduced satisfaction with the female sex assignment (Ehrhardt et al. 1968, Ehrhardt & Baker 1974, Hines et al. 2004). No evidence thus far has linked normal variability in the early hormone environment to gender dysphoria. In addition, research attempting to link 2D:4D to gender identification has produced inconsistent

findings (Kraemer et al. 2009, Schneider et al. 2006, Wallien et al. 2008), again perhaps because of the weak relationship between 2D:4D and prenatal androgen exposure.

Early Hormone Influences on Personality and Cognition

Sex differences in personality characteristics and in cognitive ability are smaller than are sex differences in children's sex-typed activities, sexual orientation, or gender identity (Hines 2010b). Nevertheless, they also have been examined for evidence of early hormonal influence.

Some personality characteristics that show sex differences relate to prenatal testosterone exposure. For instance, empathy, which is higher on average in females than in males, appears to be reduced by testosterone exposure before birth. Females with CAH show reduced empathy (Mathews et al. 2009), and testosterone measured in amniotic fluid relates negatively to empathy in both boys and girls (Chapman et al. 2006). Tendencies toward physical aggression, which are higher on average in males than in females, also relate to prenatal testosterone exposure, with prenatal testosterone exposure increasing aggression. Girls and women with CAH show increased physical aggression (Mathews et al. 2009, Pasterski et al. 2007), as do children exposed prenatally to androgenic progestins (Reinisch 1981). Not all personality dimensions that show average sex differences relate to prenatal testosterone exposure, however. For instance, the study that reported increased aggression and reduced empathy in females with CAH also considered the personality dimension of dominance/assertiveness, which is higher on average in males than in females. Despite seeing the expected sex difference in healthy controls, no difference in dominance/assertiveness was seen between females with and without CAH.

Cognitive and motor abilities that show sex differences also have been examined for influences of prenatal testosterone exposure (reviewed in Hines 2010a). One study found that

females with CAH showed more male-typical behavior in the form of increased accuracy in throwing balls and darts at targets (Hines et al. 2003), a result that was not accounted for by increased muscle strength (Collaer et al. 2009). Some studies have also found that females with CAH resemble males in showing enhanced mental rotations performance, but other studies have not corroborated these results (Hines et al. 2003). Two studies found that males with CAH show reduced performance on mental rotations or other visuo-spatial tasks (Hampson et al. 1998, Hines et al. 2003), results that had not been predicted. Several studies have also found that both males and females with CAH show impaired performance on arithmetic and mathematical tests (Baker & Ehrhardt 1974, Perlman 1973, Sinforiani et al. 1994), despite males generally being viewed as better than females at mathematics. Studies relating amniotic fluid testosterone to spatial and mathematical performance have also produced inconsistent and largely negative results (Finegan et al. 1992, Grimshaw et al. 1995). Most studies have found no alterations in individuals with CAH on tasks at which females excel, such as verbal fluency or perceptual speed, although one study suggests reduced female-typical behavior in females with CAH in the form of impaired fine motor performance (Collaer et al. 2009). Perhaps prenatal testosterone exposure has a clearer impact on motor abilities that show sex differences (e.g., targeting and fine motor performance) than on cognitive abilities assessed with paper-and-pencil tests.

Socialization and Sex Differences in Cognitive Performance

Substantial evidence supports social and cultural influences on some cognitive sex differences (see Hines 2010a for a review). For instance, sex differences on certain measures of cognitive abilities appear to have declined over time (Feingold 1988). For the SAT Mathematics, in particular, the sex ratio among those scoring at the upper extreme has declined from

13 boys to one girl in 1982 to 2.8 boys to 1 girl more recently (Halpern et al. 2007). There are also large national differences in mathematical and science performance, differences that are many fold larger than the sex difference within any nation (Mullis et al. 2008). Additionally, the magnitude of the sex difference in mathematics performance within a nation relates to the role of women. Nations where women and men are similar in regard to variables such as representation in the legislature show more equal mathematics performance (Guiso et al. 2008).

Sex-Related Psychiatric Disorders

Some psychiatric disorders are more common in one sex or the other, and testosterone could contribute here, as well. For example, prenatal testosterone exposure has been suggested to contribute to autistic spectrum conditions (ASC) (Baron-Cohen 2002) and to obsessive compulsive disorder (OCD) and Tourette Syndrome (Alexander & Peterson 2004), and to be protective against eating disorders (Culbert et al. 2008, Klump et al. 2006). For OCD and Tourette syndrome, evidence that individuals with these disorders are more male-typical in other respects, such as childhood play behavior, has been interpreted to support a link to testosterone (Alexander & Peterson 2004). For ASC (Auyeung et al. 2009a, Chapman et al. 2006, Knickmeyer et al. 2006) and for eating disorders (Culbert et al. 2008, Klump et al. 2006), behaviors in the normal range that are similar to those seen in the disorders (e.g., empathy for ASC, disordered eating for eating disorders) have been linked to prenatal androgens, although for disordered eating, some studies have failed to replicate these results (Raevuori et al. 2008). In addition, for ASC and for eating disorders, studies have not shown that variability in the early hormone environment leads to the disorder itself, as opposed to behaviors in the normal range that resemble those that characterize the disorder. For instance, although a study of females exposed to high levels of androgens prenatally, because of CAH, found increased scores on an inventory

of traits related to ASC, none of the women with CAH scored high enough to suggest a clinical diagnosis (Knickmeyer et al. 2006). The proposed link between prenatal testosterone and ASC has also been questioned by evidence indicating that both males and females with gender identity disorder, rather than females only, are at increased risk of ASC (de Vries et al. 2010) and by the larger male predominance for the less severe ASC, Asperger syndrome, than for the more severe ASC, classical autism. One possibility is that prenatal androgen exposure contributes to individual differences within the normal range in behaviors that show sex differences and that some of these resemble behaviors associated with developmental disorders, such as ASC. As a consequence, exposure to testosterone before birth, when added to other risk factors, could contribute to some individuals crossing a threshold for diagnosis. However, developmental disorders are one area in which direct genetic effects (Skuse 2006), particularly those of genes encoded in the X and Y chromosomes (Reinius et al. 2008, Reinius & Jazin 2009, Skuse 2006), may play an important role.

Sex Differences in Brain Structure and Function

There are numerous reports of sex differences in human brain structure or function (reviewed by Cahill 2009, Hines 2009). For instance, total brain volume, like body size, is larger in males than females. In addition, the amygdala is larger in males, whereas the hippocampus is larger in females (Goldstein et al. 2001). Women also show greater cortical thickness than men do in many regions (Luders et al. 2006). Perhaps in compensation for the smaller brain, women also show greater gyrfication in parts of frontal and parietal cortex and perhaps more efficient use of white matter (Gur et al. 1999). There are many reports of sex differences in the human brain, particularly in its function, and many of these are as yet unreplicated. Because males and females are routinely compared in studies, and positive results are more readily published than

negative results, some findings of neural sex differences may prove to be spurious.

In addition, although many neural sex differences have been described, few have been linked to behavioral sex differences. In fact, many differences in brain function have been noted during equivalent performance by the sexes. For instance, men and women show different patterns of asymmetry of function when performing certain phonological tasks, despite showing no sex difference in task performance (Shaywitz et al. 1995). Similarly, for men and women matched for mathematical ability, mathematical performance correlates with temporal lobe activation in men but not in women (Haier & Benbow 1995), and for women, performance on intelligence tests that do not differ by sex correlates with gray and white matter in frontal regions, whereas for men the correlation is with parietal regions (Haier et al. 2005). Indeed, neural sex differences may sometimes, or even commonly, exist to produce similar behavior in males and females, rather than to produce differences (De Vries & Sodersten 2009, McCarthy et al. 2009). Additionally, it appears that during performance of many tasks, male and female brains function similarly (Frost et al. 1999, Halari et al. 2005, Mansour et al. 1996).

Despite the many neural sex similarities and the many neural sex differences that do not relate to any behavioral sex difference, neural and behavioral sex differences have been linked in some instances. Much research in this area has focused on neural differences related to sexual orientation, particularly in men. The only finding in this area that has been independently replicated, at least as of yet, involves the third interstitial nucleus of the anterior hypothalamus (INAH-3). INAH-3 is thought to be the human homolog of the rodent SDN-POA, and four different research groups have reported that INAH-3, like the SDN-POA, is larger in males than in females (Allen et al. 1989, Byne et al. 2001, Garcia-Falgueras & Swaab, 2008, LeVay 1991). INAH-3 is also smaller (i.e., more female-typical) in homosexual than heterosexual men (Byne et al. 2001, LeVay 1991), although the number of neurons in the nucleus

INAH 1 to 4:
interstitial nuclei of
the anterior
hypothalamus,
numbers 1 to 4

appears similar for these two groups (Byne et al. 2001). The volumetric sex difference does not appear to relate to disease processes (e.g., HIV status) or to hormone use in adulthood (see Hines 2009 for discussion). Because the sex difference in SDN-POA volume in other mammals results from early testosterone exposure, differences in INAH-3 volume in humans may relate to the early hormone environment, as well. This possibility has not yet been directly investigated, however.

Heterosexual and homosexual men also differ in corpus callosum anatomy; the isthmus, in particular, is significantly larger in right-handed homosexual compared with right-handed heterosexual men (Witelson et al. 2008). Patterns of cerebral asymmetry and functional cortical connectivity have also been linked to sexual orientation in both men and women (Savic & Lindstrom 2008).

Researchers have also searched for neural correlates of gender identity disorder. One group has reported that the central subregion of the BNST (BNSTc) is smaller in women and in male-to-female transsexuals than in non-transsexual men (Zhou et al. 1995). Interpretation of this finding is complicated, however, because the sex difference in BNSTc does not appear until after puberty (Chung et al. 2002), whereas most transsexual individuals recall feeling strongly cross-gendered from early childhood. Thus, the difference in BNSTc may be the result of experience (Hines, 2009) or of the adult hormone treatment associated with changing sex (Lawrence 2009). This same research group also reported that INAH-3 is smaller and contains fewer neurons in male-to-female transsexuals than in control males (Garcia-Falgueras & Swaab 2008).

In the realm of cognitive and motor sex differences, the midsagittal area of posterior callosal regions, particularly the splenium, relates negatively to language lateralization and positively to verbal fluency in women (Hines et al. 1992b). These findings suggest a correspondence between female-typical brain structure and female-typical cognitive function, given that language lateralization is reduced

in women compared to men (McGlone, 1980, Voyer, 1996), whereas verbal fluency is greater in women than in men (Hyde & Linn 1988, Kolb & Wishaw 1985, Spreen & Strauss 1991) and posterior callosal regions tend to be larger in women than in men as well (de LaCoste-Utamsing & Holloway 1982, Witelson 1985).

Gron et al. (2000) have also described links between sex differences in brain function and navigational performance (Gron et al. 2000). In both men and women, navigating through a virtual maze, on which males perform better on average than do females, is accompanied by neural activity in the medial occipital gyri, medial and lateral superior parietal lobules, posterior cingulate and parahippocampal gyri, and the right hippocampus proper. However, women show more activity than men do in the right prefrontal cortex at Brodmann's areas 46/9, the right inferior parietal lobule, and the right superior parietal lobule, whereas men show significantly more activity than women do in the left hippocampus proper, the right parahippocampal gyrus, and the left posterior cingulate. Women with CAH have been found to perform better than healthy women on a different virtual maze task (Mueller et al. 2008), but there is, as yet, no evidence regarding neural activation in women with and without CAH while performing navigational tasks.

Effects of Experience on the Brain

Sex differences in brain structure or even function are often interpreted to imply inborn differences between males and females. This leap is inappropriate, however. Behavioral differences must be accompanied by neural differences, so the observation of a neural sex difference on its own tells us little to nothing about how the difference developed. This is true not only for differences in brain function, but also, at least in some cases, for differences in brain structure. For instance, experience can change the mammalian brain throughout the life span, and even neurogenesis in some brain regions can continue in adulthood (Juraska 1998, Maguire et al. 2006, Ming & Song 2005).

Hence, the existence of a neural sex difference, even one that relates to a behavior known to be influenced by early androgen exposure, does not prove that the hormone exposure caused the neural difference. A more direct strategy for identifying links between early hormones and the brain could be to look at neural structure or function in individuals with early hormone abnormality or in individuals for whom the early hormone environment has been measured. Although very little information of this type is available, some neural differences have been described in individuals with CAH (Hines 2009). Most notably, both males and females with CAH show decreased amygdala volume (Merke et al. 2003), and females with CAH show increased amygdala activation to negative facial emotions and, in this respect, resemble healthy males (Ernst et al. 2007). These findings fit well with expectations based on experimental work in other species because the amygdala, or some of its subregions, is larger in males than in females, contains receptors for androgen, is influenced by early manipulations of testosterone, and is involved in behaviors that show sex differences, including rough-and-tumble play and aggression (Cooke et al. 2007, Hines et al. 1992a).

Other Potential Critical Periods for Hormone Influences on Gender Development

In addition to the difference in testosterone in male and female fetuses, males and females differ in gonadal hormone levels neonatally. Shortly after birth, testosterone surges in boys (Forest et al. 1974), and estrogen surges in girls (Bidlemaier et al. 1974, 1987). The testosterone surge has been called “mini-puberty” and may play a role in development of the gonads and external genitalia in infant boys (Quigley 2002). Human brain development, particularly cortical development, continues rapidly for the first two years after birth and reacts to experience (de Graaf-Peters & Hadders-Algra 2005, Huttenlocher 2002). Thus, this early postnatal period could provide

a time when gonadal hormones and experience interact to shape the brain and behavioral propensities.

The early postnatal hormone surges, particularly the testosterone surge in boys, are a focus of current research activity. Men with anorchia (missing testes) but with normal penile development, who apparently experience normal testosterone levels prenatally, but who lack testosterone after birth, resemble controls in terms of sexual orientation, core gender identity, and questionnaire measures of personality characteristics viewed as masculine or feminine (Poomthavorn et al. 2009). Other characteristics may be influenced by the postnatal hormone surge, however. For example, men who lack the postnatal testosterone surge because they have idiopathic hypogonadotropic hypogonadism show reduced spatial abilities, and this condition is not reversed by subsequent testosterone replacement (Hier & Crowley 1982). In addition, females who do not experience the early postnatal surge of gonadal steroids because they have Turner syndrome show evidence of reduced performance on tasks at which males excel, as well as on tasks at which females excel, but not on sex-neutral tasks (Collaer et al. 2002). Evidence from healthy infants also suggests that the postnatal testosterone surge may play a role in gender development. Initial evidence suggests that testosterone during early infancy relates to infants’ visual preferences for social stimuli (Alexander et al. 2009a), to neural organization for language processing (Friederici et al. 2008), and to sex-related development of the visual system (Held et al. 1996). Although these initial reports are somewhat inconsistent and require replication, they provide intriguing glimpses through a potential new window on early gonadal hormone contributions to human gender development.

Contemporary research is also focusing on possible hormonal influences on neurobehavioral sexual differentiation at puberty. The hormonal changes of puberty produce dramatic changes in the human body, and experimental research in rodents suggests that they produce

Idiopathic hypogonadotropic hypogonadism:

involves gonadotropin deficiency and impaired gonadal steroid production after birth in affected males

Turner syndrome:

absent or imperfect second X chromosome causes ovarian regression, typically before birth, impairing or eliminating ovarian hormone production

an additional wave of neural and behavioral organization as well (Schulz et al. 2009).

Puberty is also a time of great change in human behavior, characterized by increased sexual interest and activity and the emergence of some types of behavioral problems, including higher rates of depression in females than in males (Halpern et al. 1993, Hyde et al. 2008). Evidence supports the existence of sex differences in the timing of some neural changes that accompany puberty, and these seem to parallel the earlier puberty experienced by girls compared with boys. Total cerebral volume peaks earlier in girls than in boys (at about age 10.5 years versus 14.5 years), and although both adolescent girls and adolescent boys show an inverted-U-shaped pattern of change in cortical and subcortical gray matter, the peak occurs one to two years earlier in girls than in boys (Lenroot et al. 2007). Studies have also proposed links to hormones at this time. One study found that among girls global gray matter volume related negatively to estradiol levels, but among boys the same variable related positively to testosterone levels (Peper et al. 2009). Similarly, neural sex differences in adolescent girls and boys have been found to relate to circulating testosterone levels (Neufang et al. 2009). These data are correlational, so investigators do not know if hormones, or other associated developmental processes, are the causal factors. Another study, however, suggests that testosterone may play a role in the growth of white matter in the adolescent brain. In this study, white matter increased at different rates in girls and boys, and testosterone levels and androgen receptor type interacted in relation to this sex difference. The association between male-typical brain development and testosterone levels was significantly stronger in boys with the more efficient type of androgen receptor than in boys with the less efficient type (Perrin et al. 2008). Like the investigations of possible organizational influences of hormones during neonatal development, research on puberty as an additional time of brain organization in relation to gender-linked behavior is in its early

stages, but it offers promise for understanding the dramatic behavioral changes that occur at this time of adolescent development.

CONCLUDING REMARKS

The prenatal hormone environment clearly contributes to the development of sex-related variation in human behavior and plays a role in the development of individual differences in behavior within each sex, as well as differences between the sexes. Thus, early hormone differences appear to be part of the answer to questions such as why males and females differ behaviorally and neurally, as well as why some of us are more sex-typical than others. In other species, the early hormone environment exerts its enduring effects on behavior by altering neural development. Similar neural changes are thought to underlie associations between the early hormone environment and human behavior, but the specific neural changes involved are just beginning to be identified. Many sex differences have been described in the human brain, but only a subset of these has been related to behavioral sex differences and still fewer have been linked to the early hormone environment. Steroid-sensitive regions, including regions of the hypothalamus and the amygdala, are implicated, as are interhemispheric connections, but establishing firm links between early hormones, brain development, and behavior is a primary area for future research.

Although this review has focused on hormonal influences on gender-related brain development and behavior, it has also discussed direct genetic influences that may contribute, in particular, to developmental disabilities, such as autistic spectrum conditions. In addition, the role of socialization, culture, and cognitive developmental processes in the development of behavioral differences between males and females has been noted. Although hormones contribute to behavioral sex differences, other factors contribute, as well. In addition, gender development is multidimensional, and developmental processes involved in each dimension are likely to differ somewhat. A good

example of the numerous types of factors that can influence human gender development comes from research on children's play. Here, evidence clearly shows that prenatal testosterone exposure plays a role in sex differences and individual differences, promoting male-typical toy, playmate, and activity interests. After birth, the early surges of testosterone or estrogen may be important, too, but socialization factors also gain in importance, as parents and then peers and eventually teachers encourage children to engage in gender-typed play (Fagot 1978, Langlois & Downs 1980, Pasterski et al. 2005). The child also begins to develop the understanding that he or she is male or female, and

this knowledge produces motivation to imitate the behavior of others of the same sex and to respond to information that things, such as toys or activities, are for girls or for boys by choosing the things that they have been told are for their own sex (Bussey & Bandura 1999, Martin et al. 2002, Masters et al. 1979, Perry & Bussey 1979). These social and cognitive developmental influences on children's activities could engage the same neural circuitry as underlies the effects of the early hormone environment. Thus, identifying the brain systems influenced by early androgen exposure could help elucidate systems involved in other types of influences on the same behavioral outcomes, as well.

SUMMARY POINTS

1. Human gender development begins before birth and is influenced by levels of testosterone prenatally, and perhaps neonatally. Sex-typed play in childhood relates to levels of testosterone before birth, and evidence indicates that the prenatal hormone environment also contributes to variability in sexual orientation, gender identity, and some, but not all, personality traits that differ on average for males and females.
2. Other types of influences on neurobehavioral gender development include direct genetic effects of the sex chromosomes and postnatal socialization and cognitive understanding of gender.
3. Gender development is multidimensional, and the combinations of factors that influence the many different dimensions of gender appear to differ. Early hormonal influences appear to play a larger role, for example, in children's toy preferences than they do in cognitive abilities that show sex differences, where social and cultural influences appear to be more important.

FUTURE ISSUES

1. Which neural changes can be associated with the early hormone environment, either in individuals with disorders that cause hormone abnormality or in healthy individuals for whom measures of the early hormone environment are available?
2. Does the early hormone environment contribute to the development of psychological disorders that are more common in one sex or the other?
3. Will early infancy and puberty prove to be critical periods when hormones exert permanent influences on human gendered behavior, as has been shown for prenatal development?
4. Are the neural systems associated with hormone-induced changes in behaviors that show sex differences also the systems that respond to experiential effects on the same behavioral outcomes?

DISCLOSURE STATEMENT

The author is not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

The author's research described in this article was supported in part by the United States Public Health Service (HD24542) and the Wellcome Trust (069606).

LITERATURE CITED

- Alexander GM. 2003. An evolutionary perspective of sex-typed toy preferences: pink, blue, and the brain. *Arch. Sex. Behav.* 32:7–14
- Alexander GM, Hines M. 2002. Sex differences in response to children's toys in nonhuman primates (*Cercopithecus aethiops sabaeus*). *Evol. Hum. Behav.* 23:467–79
- Alexander GM, Peterson BS. 2004. Testing the prenatal hormone hypothesis of tic-related disorders: gender identity and gender role behavior. *Dev. Psychopathol.* 16:407–20
- Alexander GM, Wilcox T, Farmer ME. 2009a. Hormone-behavior associations in early infancy. *Horm. Behav.* 56:498–502
- Alexander GM, Wilcox T, Woods R. 2009b. Sex differences in infants' visual interest in toys. *Arch. Sex. Behav.* 38:427–33
- Allen LS, Hines M, Shryne JE, Gorski RA. 1989. Two sexually dimorphic cell groups in the human brain. *J. Neurosci.* 9:497–506
- Arnold AP. 2009. The organizational-activational hypothesis as the foundation for a unified theory of sexual differentiation of all mammalian tissues. *Horm. Behav.* 55:570–78
- Auyeung B, Baron-Cohen S, Ashwin E, Knickmeyer R, Taylor K, Hackett G. 2009a. Fetal testosterone and autistic traits. *Br. J. Psychol.* 100:1–22
- Auyeung B, Baron-Cohen S, Chapman E, Knickmeyer R, Taylor K, et al. 2009b. Fetal testosterone predicts sexually differentiated childhood behavior in girls and in boys. *Psychol. Sci.* 20:144–48
- Bailey JM, Zucker KJ. 1995. Childhood sex-typed behavior and sexual orientation: a conceptual analysis and quantitative review. *Dev. Psychol.* 31:43–55
- Baker SW, Ehrhardt AA. 1974. Prenatal androgen, intelligence and cognitive sex differences. See Friedman et al. 1974, pp. 53–76
- Baron-Cohen S. 2002. The extreme male brain theory of autism. *Trends Cogn. Sci.* 6:248–54
- Berenbaum SA, Hines M. 1992. Early androgens are related to childhood sex-typed toy preferences. *Psychol. Sci.* 3:203–6
- Bidlingmaier F, Strom TM, Dörr G, Eisenmenger W, Knorr D. 1987. Estrone and estradiol concentrations in human ovaries, testes, and adrenals during the first two years of life. *J. Clin. Endocrinol. Metab.* 65:862–67
- Bidlingmaier F, Versmold H, Knorr D. 1974. Plasma estrogens in newborns and infants. In *Sexual Endocrinology of the Perinatal Period*, ed. M Forest, J Bertrand, pp. 299–314. Paris: Inserm
- Bleier R, Byne W, Siggelkow I. 1982. Cytoarchitectonic sexual dimorphisms of the medial preoptic and anterior hypothalamic areas in guinea pig, rat, hamster, and mouse. *J. Comp. Neurol.* 212:118–30
- Bussey K, Bandura A. 1999. Social cognitive theory of gender development and differentiation. *Psychol. Rev.* 106:676–713
- Byne W. 1998. The medial preoptic and anterior hypothalamic regions of the rhesus monkey: cytoarchitectonic comparison with the human and evidence for sexual dimorphism. *Brain Res.* 793:346–50
- Byne W, Tobet SA, Mattiace LA, Lasco MS, Kemether E, et al. 2001. The interstitial nuclei of the human anterior hypothalamus: an investigation of variation with sex, sexual orientation, and HIV status. *Horm. Behav.* 40:86–92
- Cahill L. 2009. Sex differences in human brain structure and function: relevance to learning and memory. See Pfaff et al. 2009, pp. 2307–15

- Campbell A, Shirley L, Heywood C. 2000. Infants' visual preference for sex-congruent babies, children, toys, and activities: a longitudinal study. *Br. J. Dev. Psychol.* 18:479–98
- Carson DJ, Okuno A, Lee PA, Stetten G, Didolkar SM, Migeon CJ. 1982. Amniotic fluid steroid levels: fetuses with adrenal hyperplasia, 46,XXY fetuses, and normal fetuses. *Am. J. Dis. Child.* 136:218–22
- Chapman E, Baron-Cohen S, Auyeung B, Knickmeyer R, Taylor K, Hackett G. 2006. Fetal testosterone and empathy: evidence from the Empathy Quotient (EQ) and the “reading the mind in the eyes” test. *Soc. Neurosci.* 1:135–48
- Chung WCJ, De Vries GJ, Swaab D. 2002. Sexual differentiation of the bed nucleus of the stria terminalis in humans may extend into adulthood. *J. Neurosci.* 22(3):1027–33
- Collaer ML, Brook CDG, Conway GS, Hindmarsh PC, Hines M. 2009. Motor development in individuals with congenital adrenal hyperplasia: strength, targeting, and fine motor skill. *Psychoneuroendocrinology* 34:249–58
- Collaer ML, Geffner M, Kaufman FR, Buckingham B, Hines M. 2002. Cognitive and behavioral characteristics of Turner syndrome: exploring a role for ovarian hormones in female sexual differentiation. *Horm. Behav.* 41:139–55
- Collaer ML, Hines M. 1995. Human behavioral sex differences: a role for gonadal hormones during early development? *Psychol. Bull.* 118:55–107
- Collaer ML, Reimers S, Manning J. 2007. Visuospatial performance on an Internet line judgment task and potential hormonal markers: sex, sexual orientation, and 2D:4D. *Arch. Sex. Behav.* 36:177–92
- Cooke BM, Stokas MR, Woolley CS. 2007. Morphological sex differences and laterality in the prepubertal medial amygdala. *J. Comp. Neurol.* 501:904–15
- Culbert KM, Breedlove SM, Burt SA, Klump KL. 2008. Prenatal hormone exposure and risk for eating disorders. *Arch. Gen. Psychiatry* 65:329–36
- de Graaf-Peters VP, Hadders-Algra M. 2005. Ontogeny of the human central nervous system: What is happening when? *Early Hum. Dev.* 82:257–66
- de Lacoste-Utamsing C, Holloway RL. 1982. Sexual dimorphism in the human corpus callosum. *Science* 216:1431–32
- Dessens AB, Slijper FME, Drop SLS. 2005. Gender dysphoria and gender change in chromosomal females with congenital adrenal hyperplasia. *Arch. Sex. Behav.* 34:389–97
- de Vries ALC, Noens ILJ, Cohen-Kettenis PT, Berckelaer-Onnes IA, Doreleijers TA. 2010. Autism spectrum disorders in gender dysphoric children and adolescents. *J. Autism Dev. Disord.* 40:930–36
- De Vries GJ, Sodersten P. 2009. Sex differences in the brain: the relation between structure and function. *Horm. Behav.* 55:589–96
- Dittmann RW, Kappes MH, Kappes ME, Börger D, Stegner H, et al. 1990. Congenital adrenal hyperplasia I: gender-related behavior and attitudes in female patients and sisters. *Psychoneuroendocrinology* 15:401–20
- Dohler KD, Coquelin A, Davis F, Hines M, Shryne JE, Gorski RA. 1984. Pre- and postnatal influence of testosterone propionate and diethylstilbestrol on differentiation of the sexually dimorphic nucleus of the preoptic area in male and female rats. *Brain Res.* 302:291–95
- Ehrhardt AA, Baker SW. 1974. Fetal androgens, human central nervous system differentiation, and behavior sex differences. See Friedman et al. 1974, pp. 33–52
- Ehrhardt AA, Epstein R, Money J. 1968. Fetal androgens and female gender identity in the early-treated adrenogenital syndrome. *Johns Hopkins Med. J.* 122:160–67
- Ehrhardt AA, Grisanti GC, Meyer-Bahlburg HFL. 1977. Prenatal exposure to medroxyprogesterone acetate (MPA) in girls. *Psychoneuroendocrinology* 2:391–98
- Ehrhardt AA, Money J. 1967. Progestin-induced hermaphroditism: IQ and psychosexual identity in a study of ten girls. *J. Sex Res.* 3:83–100
- Ernst M, Maheu FS, Schroth E, Hardin J, Golan LS, et al. 2007. Amygdala function in adolescents with congenital adrenal hyperplasia: a model for the study of early steroid abnormalities. *Neuropsychologia* 45:2104–13
- Fagot BI. 1978. The influence of sex of child on parental reactions to toddler children. *Child Dev.* 49:459–65
- Fausto-Sterling A. 1992. *Myths of Gender*. New York: Basic Books

- Feingold A. 1988. Cognitive gender differences are disappearing. *Am. Psychol.* 43:95–103
- Finegan JK, Niccols GA, Sitarenios G. 1992. Relations between prenatal testosterone levels and cognitive abilities at 4 years. *Dev. Psychol.* 28:1075–89
- Forest MG, Sizonenko PC, Cathiard AM, Bertrand J. 1974. Hypophyso-gonadal function in humans during the first year of life. I. Evidence for testicular activity in early infancy. *J. Clin. Invest.* 53:819–28
- Friederici AD, Pannekamp A, Partsch C-J, Ulmen U, Oehler K et al. 2008. Sex hormone testosterone affects language organization in the infant brain. *Neuroreport* 19:283–86
- Friedman RC, Richart RN, Vande Wiele RL, eds. 1974. *Sex Differences in Behavior*. New York: Wiley
- Frost JA, Binder JR, Springer JA, Hammeke TA, Bellgowan SF, et al. 1999. Language processing is strongly left lateralized in both sexes—evidence from functional MRI. *Brain* 122:199–208
- Garcia-Falgueras A, Swaab DF. 2008. A sex difference in the hypothalamic uncinate nucleus: relationship to gender identity. *Brain* 131:3132–46
- Ginsburg KR, Committee on Communications, Committee on Psychosocial Aspects of Child and Family Health. 2007. The importance of play in promoting healthy child development and maintaining strong parent-child bonds. *Pediatrics* 119:182–91
- Goldstein JM, Seidman LJ, Horton NJ, Makris N, Kennedy DN, et al. 2001. Normal sexual dimorphism of the adult human brain assessed by in vivo magnetic resonance imaging. *Cereb. Cortex* 11:490–97
- Gorski RA, Gordon JH, Shryne JE, Southam AM. 1978. Evidence for a morphological sex difference within the medial preoptic area of the rat brain. *Brain Res.* 148:333–46
- Gorski RA, Harlan RE, Jacobson CD, Shryne JE, Southam AM. 1980. Evidence for the existence of a sexually dimorphic nucleus in the preoptic area of the rat. *J. Comp. Neurol.* 193:529–39
- Green R. 1985. Gender identity in childhood and later sexual orientation. *Am. J. Psychiatry* 142:339–41
- Grimbos T, Dawood K, Burriss RP, Zucker KJ, Puts DA. 2010. Sexual orientation and the second to fourth finger length ratio: a meta-analysis in men and women. *Behav. Neurosci.* 124:278–87
- Grimshaw GM, Sitarenios G, Finegan JK. 1995. Mental rotation at 7 years: relations with prenatal testosterone levels and spatial play experiences. *Brain Cogn.* 29:85–100
- Gron G, Wunderlich AP, Spitzer M, Tomczak R, Riepe MW. 2000. Brain activation during human navigation: gender-different neural networks as substrate of performance. *Nat. Neurosci.* 3:404–8
- Guiso L, Monte F, Sapienza P, Zingales L. 2008. Culture, gender and math. *Science* 320:1164–65
- Gur RC, Turetsky BI, Matsui M, Yan M, Bilker W, et al. 1999. Sex differences in gray and white matter in healthy young adults: correlations with cognitive performance. *J. Neurosci.* 19:4065–72
- Haier RJ, Benbow CP. 1995. Sex differences and lateralization in temporal lobe glucose metabolism during mathematical reasoning. *Dev. Neuropsychol.* 11:405–14
- Haier RJ, Jung RE, Yeo RA, Head K, Alkire MT. 2005. The neuroanatomy of general intelligence: Sex matters. *Neuroimage* 25:320–27
- Halari R, Sharma T, Hines M, Andrew C, Simmons A, Kumari V. 2005. Comparable fMRI activity with differential behavioural performance on mental rotation and overt verbal fluency tasks in healthy men and women. *Exp. Brain Res.* 169:1–14
- Hall CM, Jones JA, Meyer-Bahlburg HFL, Dolezal C, Coleman M, et al. 2004. Behavioral and physical masculinization are related to genotype in girls with congenital adrenal hyperplasia. *J. Clin. Endocrinol. Metab.* 89:419–24
- Halpern CT, Udry JR, Campbell B, Suchindran C. 1993. Testosterone and pubertal development as predictors of sexual activity: a panel analysis of adolescent males. *Psychosom. Med.* 55:436–47
- Halpern DF, Benbow CP, Geary DC, Gur RC, Hyde JS, Gernsbacher MA. 2007. The science of sex differences in science and mathematics. *Psychol. Sci. Public Interest* 8:1–51
- Hampson E, Rovet J, Altmann D. 1998. Spatial reasoning in children with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Dev. Neuropsychol.* 14(2):299–320
- Hassett JM, Siebert ER, Wallen K. 2008. Sex differences in rhesus monkey toy preferences parallel those of children. *Horm. Behav.* 54:359–64
- Held R, Thorn F, Gwiazda J, Bauer J. 1996. Development of binocularity and its sexual differentiation. In *Infant Vision*, ed. F Vital-Durant, J Atkinson, OJ Braddick, pp. 265–74. New York: Oxford Univ. Press
- Hier DB, Crowley WF. 1982. Spatial ability in androgen-deficient men. *N. Engl. J. Med.* 306:1202–5

- Hines M. 2004. *Brain Gender*. New York: Oxford Univ. Press
- Hines M. 2009. Gonadal hormones and sexual differentiation of human brain and behavior. See Pfaff et al. 2009, pp. 1869–909
- Hines M. 2010a. Gendered behavior across the lifespan. In *Life-Span Development*, Vol. 2, ed. RM Lerner, ME Lamb, A Freund, pp. 341–78. New York: Wiley
- Hines M. 2010b. Sex-related variation in human behavior and the brain. *Trends Cogn. Sci.* 14:448–56
- Hines M, Allen LS, Gorski RA. 1992a. Sex differences in subregions of the medial nucleus of the amygdala and the bed nucleus of the stria terminalis of the rat. *Brain Res.* 579:321–26
- Hines M, Alsum P, Roy M, Gorski RA, Goy RW. 1987. Estrogenic contributions to sexual differentiation in the female guinea pig: influences of diethylstilbestrol and tamoxifen on neural, behavioral and ovarian development. *Horm. Behav.* 21:402–17
- Hines M, Brook C, Conway GS. 2004. Androgen and psychosexual development: core gender identity, sexual orientation and recalled childhood gender role behavior in women and men with congenital adrenal hyperplasia (CAH). *J. Sex Res.* 41:75–81
- Hines M, Chiu L, McAdams LA, Bentler PM, Lipcamon J. 1992b. Cognition and the corpus callosum: verbal fluency, visuospatial ability and language lateralization related to midsagittal surface areas of callosal subregions. *Behav. Neurosci.* 106:3–14
- Hines M, Fane BA, Pasterski VL, Mathews GA, Conway GS, Brook C. 2003. Spatial abilities following prenatal androgen abnormality: targeting and mental rotations performance in individuals with congenital adrenal hyperplasia (CAH). *Psychoneuroendocrinology* 28:1010–26
- Hines M, Golombok S, Rust J, Johnston K, Golding J, ALSPAC Study Team. 2002. Testosterone during pregnancy and childhood gender role behavior: a longitudinal population study. *Child Dev.* 73:1678–87
- Huttenlocher PR. 2002. *Neural Plasticity*. Cambridge, MA: Harvard Univ. Press
- Hyde JS, Mezulis AH, Abramson LY. 2008. The ABCs of depression: integrating affective, biological, and cognitive models to explain the emergence of the gender difference in depression. *Psychol. Rev.* 115:291–313
- Hyde JS, Linn MC. 1988. Gender differences in verbal ability: a meta-analysis. *Psychol. Bull.* 104:53–69
- Ito S, Murakami S, Yamanouchi K, Arai Y. 1986. Perinatal androgen exposure decreases the size of the sexually dimorphic medial preoptic nucleus in the rat. *Proc. Japan Acad.* 62(B):408–11
- Jacobson CD, Csernus VJ, Shryne JE, Gorski RA. 1981. The influence of gonadectomy, androgen exposure, or a gonadal graft in the neonatal rat on the volume of the sexually dimorphic nucleus of the preoptic area. *J. Neurosci.* 1:1142–47
- Jadva V, Golombok S, Hines M. 2010. Infants' preferences for toys, colors and shapes. *Arch. Sex. Behav.* 39:1261–73
- Jordan-Young RM. 2010. *Brainstorm: The Flaws in the Science of Sex Differences*. Cambridge, MA: Harvard Univ. Press
- Juraska JM. 1998. Neural plasticity and the development of sex differences. *Annu. Rev. Sex Res.* 9:20–38
- Klump KL, Gobrogge KL, Perkins PS, Thorne D, Sisk CL, Breedlove SM. 2006. Preliminary evidence that gonadal hormones organize and activate disordered eating. *Psychol. Med.* 36:539–46
- Knickmeyer R, Baron-Cohen S, Fane B, Wheelwright S, Mathews GA, et al. 2006. Androgen and autistic traits: a study of individuals with congenital adrenal hyperplasia. *Horm. Behav.* 50:148–53
- Kolb B, Whishaw IQ. 1985. *Fundamentals of Human Neuropsychology*. New York: W.H. Freeman and Co.
- Kraemer B, Noll T, Delsignore A, Milos G, Schnyder U, Hepp U. 2009. Finger length ratio (2D:4D) in adults with gender identity disorder. *Arch. Sex. Behav.* 38:359–63
- Langlois JH, Downs AC. 1980. Mothers, fathers and peers as socialization agents of sex-typed play behaviors in young children. *Child Dev.* 51:1217–47
- Lawrence AA. 2009. Parallels between gender identity disorder and body integrity identity disorder: a review and update. In *Body Integrity Identity Disorder: Psychological, Neurobiological, Ethical, and Legal Aspects*, ed. A Stirn, A Thiel, S Oddo, pp. 154–72. Lengerich, Germ.: Pabst
- Lenroot RK, Gogtay N, Greenstein DK, Wells EM, Wallace GL, et al. 2007. Sexual dimorphism of brain developmental trajectories during childhood and adolescence. *Neuroimage* 36:1065–73

- LeVay S. 1991. A difference in hypothalamic structure between heterosexual and homosexual men. *Science* 253:1034–37
- Luders E, Narr KI, Thompson PM, Rex DE, Woods RP, et al. 2006. Gender effects on cortical thickness and the influence of scaling. *Hum. Brain Mapp.* 27:314–24
- Maguire EA, Woollett K, Spiers HJ. 2006. London taxi drivers and bus drivers: a structural MRI and neuropsychological analysis. *Hippocampus* 16:1091–101
- Mansour CS, Haier RJ, Buchsbaum MS. 1996. Gender comparisons of cerebral glucose metabolic rate in healthy adults during a cognitive task. *Pers. Individ. Differ.* 20:183–91
- Martin CL, Ruble DN, Szkrybalo J. 2002. Cognitive theories of early gender development. *Psychol. Bull.* 128:903–33
- Masters JC, Ford ME, Arend R, Grotevant HD, Clark LV. 1979. Modeling and labelling as integrated determinants of children's sex-typed imitative behavior. *Child Dev.* 50:364–71
- Mathews GA, Fane BA, Conway GS, Brook C, Hines M. 2009. Personality and congenital adrenal hyperplasia: possible effects of prenatal androgen exposure. *Horm. Behav.* 55:285–91
- McCarthy MM, De Vries GJ, Forger NG. 2009. Sexual differentiation of the brain: mode, mechanisms, and meaning. See Pfaff et al. 2009, pp. 1707–44
- McGlone J. 1980. Sex differences in human brain asymmetry: a critical survey. *Behav. Brain Sci.* 3:215–63
- Merke DP, Fields JD, Keil MF, Vaituzis AC, Chrousos GP, Giedd JN. 2003. Children with classic congenital adrenal hyperplasia have decreased amygdala volume: potential prenatal and postnatal hormonal effects. *J. Clin. Endocrinol. Metab.* 88(4):1760–65
- Meyer-Bahlburg HFL, Dolezal C, Baker SW, New MI. 2008. Sexual orientation in women with classical or non-classical congenital adrenal hyperplasia as a function of degree of prenatal androgen excess. *Arch. Sex. Behav.* 37:85–99
- Ming G, Song H. 2005. Adult neurogenesis in the mammalian central nervous system. *Annu. Rev. Neurosci.* 28:223–50
- Mueller SC, Temple V, Oh E, VanRyzin C, Williams A, et al. 2008. Early androgen exposure modulates spatial cognition in congenital adrenal hyperplasia (CAH). *Psychoneuroendocrinology* 33:973–80
- Mullis IVS, Martin MO, Foy P. 2008. *TIMSS 2007 International Maths Report*. Boston: Int. Assoc. Eval. Educ. Achiev.
- Neufang S, Specht K, Hausmann M, Gunturkun O, Herpertz-Dahlmann B, et al. 2009. Sex differences and the impact of steroid hormones on the developing human brain. *Cereb. Cortex* 19:464–73
- Nordenstrom A, Servin A, Bohlin G, Larsson A, Wedell A. 2002. Sex-typed toy play behavior correlates with the degree of prenatal androgen exposure assessed by CYP21 genotype in girls with congenital adrenal hyperplasia. *J. Clin. Endocrinol. Metab.* 87:5119–24
- Pasterski VL, Geffner ME, Brain C, Hindmarsh P, Brook C, Hines M. 2005. Prenatal hormones and postnatal socialization by parents as determinants of male-typical toy play in girls with congenital adrenal hyperplasia. *Child Dev.* 76:264–78
- Pasterski VL, Hindmarsh P, Geffner M, Brook C, Brain C, Hines M. 2007. Increased aggression and activity level in 3- to 11-year-old girls with congenital adrenal hyperplasia (CAH). *Horm. Behav.* 52:368–74
- Peper JS, Brouwer RM, Schnack HG, van Baal GC, van Leeuwen M, Van Den Berg SM, et al. 2009. Sex steroids and brain structure in pubertal boys and girls. *Psychoneuroendocrinology* 34:332–42
- Perlman SM. 1973. Cognitive abilities of children with hormone abnormalities: screening by psychoeducational tests. *J. Learn. Disabil.* 6:21–29
- Perrin JS, Herve P-Y, Leonard G, Perron M, Pike GB, et al. 2008. Growth of white matter in the adolescent brain: role of testosterone and androgen receptor. *J. Neurosci.* 28:9519–24
- Perry DG, Bussey K. 1979. The social learning theory of sex difference: Imitation is alive and well. *J. Pers. Soc. Psychol.* 37:1699–712
- Pfaff DW, Arnold AP, Etgen AM, Fahrbach SE, Rubin RT, eds. 2009. *Hormones, Brain and Behavior*. San Diego: Academic
- Phoenix CH, Goy RW, Gerall AA, Young WC. 1959. Organizing action of prenatally administered testosterone propionate on the tissues mediating mating behavior in the female guinea pig. *Endocrinology* 65:163–96

- Piaget J. 1970. *Science of Education and the Psychology of the Child*. New York: Orion
- Poomthavorn P, Stargatt R, Zacharin M. 2009. Psychosexual and psychosocial functions of anorchid young adults. *J. Clin. Endocrinol. Metab.* 94:2502–5
- Quigley CA. 2002. The postnatal gonadotrophin and sex steroid surge—insights from the androgen insensitivity syndrome. *Clin. Endocrinol. Metab.* 87(1):24–28
- Raevuori A, Kaprio J, Hoek HW, Sihvola E, Rissanen A, Keski-Rahkonen A. 2008. Anorexia and bulimia nervosa in same-sex and opposite-sex twins: lack of an association with twin type in a nationwide study of Finnish twins. *Am. J. Psychiatry* 165:1604–10
- Reinisch JM. 1981. Prenatal exposure to synthetic progestins increases potential for aggression in humans. *Science* 211:1171–73
- Reinius B, Jazin E. 2009. Prenatal sex differences in the human brain. *Mol. Psychiatry* 14:988–89
- Reinius B, Saetre P, Leonard JA, Blekhan R, Merino-Martinez R, et al. 2008. An evolutionarily conserved sexual signature in the primate brain. *PLoS Genet.* 4:e1000100
- Reyes FI, Winter JSD, Faiman C. 1973. Studies on human sexual development. I. Fetal gonadal and adrenal sex steroids. *J. Clin. Endocrinol. Metab.* 37:74–78
- Roselli CE, Larkin K, Resko JA, Stellflug JN, Stromshak F. 2004. The volume of a sexually dimorphic nucleus in the ovine medial preoptic area/anterior hypothalamus varies with sexual partner preference. *Endocrinology* 145(2):478–83
- Ruble DN, Martin CL, Berenbaum SA. 2006. Gender development. In *Handbook of Child Psychology*, Vol. 3: *Social, Emotional and Personality Development*, ed. W Damon, R M Lerner, N Eisenberg, pp. 858–932. New York: Wiley
- Savic I, Lindstrom P. 2008. PET and MRI show differences in cerebral asymmetry and functional connectivity between homo- and heterosexual subjects. *Proc. Natl. Acad. Sci. USA* 105:9403–8
- Schneider HJ, Pickel J, Stalla GK. 2006. Typical female 2nd–4th finger length (2D:4D) ratios in male-to-female transsexuals—possible implications for prenatal androgen exposure. *Psychoneuroendocrinology* 31:265–69
- Schulz KM, Molenda-Figueira HA, Sisk CL. 2009. Back to the future: the organizational-activational hypothesis adapted to puberty and adolescence. *Horm. Behav.* 55:597–604
- Serbin LA, Poulin-Dubois D, Colbourne KA, Sen MG, Eichstedt JA. 2001. Gender stereotyping in infancy: visual preferences for and knowledge of gender-stereotyped toys in the second year. *Int. J. Behav. Dev.* 25:7–15
- Shaywitz BA, Shaywitz SE, Pugh KR, Constable RT, Skudlarski P, et al. 1995. Sex differences in the functional organization of the brain for language. *Nature* 373:607–9
- Simerly RB, Zee MC, Pendleton JW, Lubahn DB, Korach KS. 1997. Estrogen receptor-dependent sexual differentiation of dopaminergic neurons in the preoptic region of the mouse. *Proc. Natl. Acad. Sci. USA* 94:14077–82
- Sinforiani E, Livieri C, Mauri M, Bisio P, Sibilla L, et al. 1994. Cognitive and neuroradiological findings in congenital adrenal hyperplasia. *Psychoneuroendocrinology* 19:55–64
- Skuse DH. 2006. Sexual dimorphism in cognition and behaviour: the role of X-linked genes. *Eur. J. Endocrinol.* 155:S99–106
- Slijper FME, Drop SLS, Molenaar JC, de Muinck Keizer-Schrama SMPF. 1998. Long-term psychological evaluation of intersex children. *Arch. Sex. Behav.* 27:125–44
- Spreen O, Strauss E. 1991. *A Compendium of Neuropsychological Tests*. New York: Oxford Univ. Press
- Sumida H, Nishizuka M, Kano Y, Arai Y. 1993. Sex differences in the anteroventral periventricular nucleus of the preoptic area and in the related effects of androgen in prenatal rats. *Neurosci. Lett.* 151:41–44
- Tobet SA, Zahniser DJ, Baum MJ. 1986. Differentiation in male ferrets of a sexually dimorphic nucleus of the preoptic/anterior hypothalamic area requires prenatal estrogen. *Neuroendocrinology* 44:299–308
- Ulibarri C, Yahr P. 1988. Role of neonatal androgens in sexual differentiation of brain structure, scent marking and gonadotropin secretion in gerbils. *Behav. Neural Biol.* 49:27–44
- Voyer D. 1996. On the magnitude of laterality effects and sex differences in functional lateralities. *Laterality* 1:51–83
- Vygotsky LS. 1976. Play and its role in the mental development of the child. *Sov. Psychol.* 5:6–18
- Wallien MSC, Zucker KJ, Steensma TD, Cohen-Kettenis PT. 2008. 2D:4D finger-length ratios in children and adults with gender identity disorder. *Horm. Behav.* 54:450–54

- Wilson JD, George FW, Griffin JE. 1981. The hormonal control of sexual development. *Science* 211:1278–84
- Witelson SF. 1989. Hand and sex differences in the isthmus and genu of the human corpus callosum: a postmortem morphological study. *Brain* 112:799–835
- Witelson SF, Kigar DL, Scamvougeras A, Kideckel DM, Buck B, et al. 2008. Corpus callosum anatomy in right-handed homosexual and heterosexual men. *Arch. Sex. Behav.* 37:857–63
- Zhou J, Hofman MA, Gooren LJG, Swaab DF. 1995. A sex difference in the human brain and its relation to transsexuality. *Nature* 378:68–70



Contents

Collision Detection as a Model for Sensory-Motor Integration <i>Haleh Fotowat and Fabrizio Gabbiani</i>	1
Myelin Regeneration: A Recapitulation of Development? <i>Stephen P.J. Fancy, Jonah R. Chan, Sergio E. Baranzini, Robin J.M. Franklin, and David H. Rowitch</i>	21
Neural Representations for Object Perception: Structure, Category, and Adaptive Coding <i>Zoe Kourtzi and Charles E. Connor</i>	45
Gender Development and the Human Brain <i>Melissa Hines</i>	69
Too Many Cooks? Intrinsic and Synaptic Homeostatic Mechanisms in Cortical Circuit Refinement <i>Gina Turrigiano</i>	89
Reward, Addiction, Withdrawal to Nicotine <i>Mariella De Biasi and John A. Dani</i>	105
Neuronal Intrinsic Mechanisms of Axon Regeneration <i>Kai Liu, Andrea Tedeschi, Kevin Kyungsuk Park, and Zbigang He</i>	131
Transcriptional Control of the Terminal Fate of Monoaminergic Neurons <i>Nuria Flames and Oliver Hobert</i>	153
Amyloid Precursor Protein Processing and Alzheimer's Disease <i>Richard J. O'Brien and Philip C. Wong</i>	185
Motor Functions of the Superior Colliculus <i>Neeraj J. Gandhi and Husam A. Katnani</i>	205
Olfactory Maps in the Brain <i>Venkatesh N. Murthy</i>	233
The Cognitive Neuroscience of Human Memory Since H.M. <i>Larry R. Squire and John T. Wixted</i>	259
Deep Brain Stimulation for Psychiatric Disorders <i>Paul E. Holtzheimer and Helen S. Mayberg</i>	289

Three-Dimensional Transformations for Goal-Directed Action <i>J. Douglas Crawford, Denise Y.P. Henriques, and W. Pieter Medendorp</i>	309
Neurobiology of Economic Choice: A Good-Based Model <i>Camillo Padoa-Schioppa</i>	333
The Extraction of 3D Shape in the Visual System of Human and Nonhuman Primates <i>Guy A. Orban</i>	361
The Development and Application of Optogenetics <i>Lief Fenno, Ofer Yizhar, and Karl Deisseroth</i>	389
Recovery of Locomotion After Spinal Cord Injury: Some Facts and Mechanisms <i>Serge Rossignol and Alain Frigon</i>	413
Modulation of Striatal Projection Systems by Dopamine <i>Charles R. Gerfen and D. James Surmeier</i>	441
How Is the Olfactory Map Formed and Interpreted in the Mammalian Brain? <i>Kensaku Mori and Hitoshi Sakano</i>	467
Vestibular Hair Cells and Afferents: Two Channels for Head Motion Signals <i>Ruth Anne Eatock and Jocelyn E. Songer</i>	501
Mechanisms of Inhibition within the Telencephalon: “Where the Wild Things Are” <i>Gord Fishell and Bernardo Rudy</i>	535
Spatial Neglect and Attention Networks <i>Maurizio Corbetta and Gordon L. Shulman</i>	569
General Anesthesia and Altered States of Arousal: A Systems Neuroscience Analysis <i>Emery N. Brown, Patrick L. Purdon, and Christa J. Van Dort</i>	601

Indexes

Cumulative Index of Contributing Authors, Volumes 25–34	629
Cumulative Index of Chapter Titles, Volumes 25–34	633

Errata

An online log of corrections to *Annual Review of Neuroscience* articles may be found at <http://neuro.annualreviews.org/>