

If you had less than a year to live, would you want to know? A seven-country European population survey of public preferences for disclosure of poor prognosis

R. Harding^{1*}, V. Simms^{1,2}, N. Calanzani¹, I. J. Higginson¹, S. Hall¹, M. Gysels³, A. Meñaca³, C. Bausewein^{1,4}, L. Deliens^{5,6}, P. Ferreira⁷, F. Toscani⁸, B. A. Daveson¹, L. Ceulemans⁹, B. Gomes¹ and on behalf of PRISMA

¹King's College London, Cicely Saunders Institute, Department of Palliative Care, Policy & Rehabilitation, London SE5 9PJ, UK

²London School of Hygiene and Tropical Medicine, London, UK

³Barcelona Centre for International Health Research (CRESB – Hospital Clínic), Universitat de Barcelona, Barcelona, Spain

⁴Deutsche Gesellschaft für Palliativmedizin, Berlin, Germany

⁵End-of-Life Care Research Group, Ghent University & Vrije Universiteit Brussel, Brussels, Belgium

⁶Department of Public and Occupational Health, EMGO Institute for Health and Care Research, Palliative Care Center of Expertise, VU University Medical Center, Amsterdam, the Netherlands

⁷Centre for Health Studies and Research, University of Coimbra (CEISUC), Coimbra, Portugal

⁸Istituto di Ricerca in Medicina Palliativa, Fondazione Lino Maestroni – ONLUS, Cremona, Italy

⁹University of Antwerp, Antwerp, Belgium

*Correspondence to:

King's College London, Cicely Saunders Institute, Department of Palliative Care, Policy & Rehabilitation, Bessemer Road, London SE5 9PJ, UK. E-mail: richard.harding@kcl.ac.uk

Abstract

Objective: With increasing European cancer deaths, clinicians must manage information regarding poor prognosis. This study aimed to determine European citizens' preferences, within a scenario of serious illness such as cancer with less than a year to live, for information disclosure regarding poor prognosis, the likely symptoms and problems, and the care options available, to measure variations between countries and to identify factors associated with preferences.

Methods: A population-based cross-national telephone survey using random digit dialling in seven countries was conducted.

Results: Among 9344 respondents, data revealed an international preference (73.9%) to always be informed in the scenario of having a serious illness such as cancer with less than a year to live. This varied from 67.6% in Italy to 80.7% in Flanders. A minority (21.1%) did not want such information unless they ask, or at all. People younger than 70 years (OR 0.72, 95% CI 0.62–0.83, $p < 0.001$), men (OR 1.23, 95% CI 1.10–1.37, $p < 0.001$), those with experience of illness (OR = 1.20, 95% CI 1.01–1.43, $p < 0.05$) and with more education (OR = 1.20, 95% CI 1.09–1.32, $p < 0.001$) were more likely to want to know of limited time left.

Conclusions: The models confirmed the influence of four factors in more than one country (age, gender, education and most concerning problem) and added 11 country-specific factors to which national policies and clinical practice should respond. These findings confirm a majority public preference to be informed in a scenario of poor prognosis. Policy clinical practice should facilitate elucidation and delivery of preferences. Evidence for effective communication skills-building interventions for clinicians is required.

Copyright © 2013 John Wiley & Sons, Ltd.

Received: 30 August 2012

Revised: 17 December 2012

Accepted: 18 February 2013

Introduction

With an ageing European population and an annual 1.7 million cancer deaths in Europe [1], clinicians increasingly need to manage information-giving in the face of poor prognosis.

Information is a primary need in advanced cancer [2]. Although being able to control the content and extent of information is important to care recipients in the context of cancer [3], it is difficult to formulate and pose sensitive questions to clinicians without being prompted [4]. Clinicians also feel unsure about when is the best time to provide information and how much information is wanted by their patients [5], they often do not share prognostic information [6], and wait to be asked rather than taking the initiative to initiate conversations [3]. As a result, the condition is misunderstood by those affected [7,8], which may increase

anxiety, reduce the potential for advance care planning and increase the likelihood of unplanned admissions.

The evidence is equivocal on how much information is wanted in advanced disease [8]. With increased public access via the internet to health-related information of variable quality, it is essential that accurate information specific to an individual's situation is available when wanted. Information-giving is an essential aspect of clinical care, and understanding public preferences for information in advanced disease is necessary to train and support clinicians in responding appropriately. From a public health perspective, it is also important to educate the public concerning the importance of communicating their preferences to their health care providers. As more people die at an advanced age, clinicians will be increasingly confronted with patients that lack mental capacity; hence, early effective information-giving becomes more important.

European medical cultures may be changing from a historical point of concealment of prognostic information justified by beneficence [9]. Differences persist within and between European populations, with recent evidence revealing that in addition to age, personal values and circumstance, country of residence also influences preferred place of death [10]. However, little is known about preferences for prognostic information among European populations and the variations between them. Moreover, little is known on how exposure to illness, caregiving and age may affect preferences. Evidence from Europe is needed to appraise and inform national and international care practices. The present study is the first to cross-nationally determine variations in people's preferences for information regarding prognosis in a scenario of advanced cancer.

This study aimed to determine the preferences of European citizens for disclosure of information regarding a poor prognosis, the likely symptoms and problems, and the care options available. The study further aimed to measure variations between countries and to identify factors associated with preferences.

Methods

Design

A population-based cross-national telephone survey in seven European countries was conducted. The details of the survey are described elsewhere, and it was undertaken as part of a pan-European clinical academic programme to enhance end-of-life cancer care for European citizens [10,11].

Questionnaire

Development

We took a multi-method approach to questionnaire development to enhance validity and comparability. This included the following: (i) review of studies and questions on end-of-life care (EoLC) preferences and priorities; (ii) review of cross-national social surveys; and (iii) three consultation rounds with 27 EoLC expert researchers and clinicians.

Content

The questionnaire included 28 items on preferences, personal values related to EoLC, experience with illness, death and dying, general health and sociodemographics. Preferences were asked in a scenario imagining 'a situation of serious illness, for example cancer, with less than one year to live'. Most demographic items were adapted from the European Social Survey Round 4 (2008) [12].

The first three survey questions are the outcomes reported in this paper: (i) Would you like to be informed that you had limited time left? (ii) Would you like to be informed about what symptoms and problems you were likely to experience? and (iii) Would you like to be informed about the options available for care and how they might affect you? These options might be services available, places where you could be looked after, treatments and medication.

Possible responses were 'Yes, always', 'Yes, but only if you ask about it', 'No' and 'You don't know'. The three

outcomes are referred to as (i) 'time left', (ii) 'symptoms' and (iii) 'care options'.

Translation

We undertook systematic and culture-sensitive translations of the questionnaire into the countries' dominant language following European Organisation for Research and Treatment in Cancer translation procedures [13]. This involved forward independent translations by two native speakers with EoLC knowledge, backward professional translation and harmonisation of all versions.

Testing

The resulting questionnaire (in supplementary data) was piloted using cognitive interviewing to check how items are interpreted with 30 volunteers in England and Germany [14].

Setting

The survey covered Flanders (the Dutch-speaking part of Belgium) and all regions in England, Germany, Italy, Netherlands, Spain and Portugal. These countries have contrasting cultural, political contexts and sociodemographic profiles [15–18].

Participants

Individuals aged ≥ 16 years residing in a household were invited to participate in a computer-assisted telephone interview by selection of households using random digit dialling. Once an eligible participant was identified, no substitution was allowed. Exclusion criteria were as follows: incapacity to hear or understand the information and provide informed consent (assessed by interviewers) and language skills that hindered survey participation (using the country's dominant language).

To be able to detect previous age differences in preferences for home death in England between those aged 16–44 years (60%) and those aged 45 years and older (51%) [19], we needed a minimum of 1278 completed interviews (power 0.90, significance 0.05). Modelling calculations to other countries resulted in overall minimum needed interviews being 8946. No oversampling, strata or quota were applied.

Procedures

The lead academic team trained 149 interviewers (95% native speakers) from two commercial research agencies with experience in telephone surveys on social and health issues on how to administer the questionnaire. Interviews were conducted from May to December 2010 with at least four call attempts (at least one after 6pm). Interviewers entered answers into a database with missing data checks at entry; 10% of interviews were checked by *in situ* supervisors.

Ethical approval

The study was approved by the research ethics committee of the lead academic centre, King's College London (ref. BDM/08/09-48). Local research ethics approvals or

European citizens' preference to be told of poor prognosis

notifications to national data protection agencies as applicable were obtained in all countries.

Analysis

The analysis was conducted by VS using Stata 10.1. Participant characteristics were presented by country for sociodemographic, experiential and opinion variables. For each of the three outcomes, the proportion of each possible response was calculated overall and by country.

The primary outcome (wish to know that there was limited time left) was converted into a binary score (yes always = 1 vs all other options = 0). Prevalence of wish to know was presented by covariate and country for 17 independent variables identified in the literature as possibly associated with information and decision-making preferences at the end of life. The independent variables were grouped as follows. Sociodemographic: gender (binary); age band (ordinal); education (ordinal: up to age 16 years, age 16–18 years and post 18 years); living area (ordinal: city, suburb, town and village/rural); general health (ordinal: 5-point scale); marital status (categorical); living in country of birth (binary); length of time in country (ordinal); financial hardship (ordinal: four categories); and living alone (binary). Illness experience in the past 5 years: serious illness of a close/relative friend (binary), death of a close relative/friend (binary), diagnosis of serious illness (binary) and ever cared for a relative/friend in the last few months of their life (binary). Opinions: preference for quality or quantity of life (ordinal: three categories), most concerning problem (binary: physical/psychological vs social) and prefer to die at home (binary).

The significance of variations in preferences by each of the 17 independent variables within and across countries was determined using χ^2 tests or Fisher's Exact Test when required for categorical data, Mann–Whitney U tests for ordinal data and t -tests for continuous data. Any variable associated with the outcome at a significance level of $p < 0.1$ for all countries combined, with shared directionality in a majority of countries, was carried forward into multivariate analysis.

Time-series logit analysis was conducted to identify influencing factors in each scenario, with country designated the panel variable. This analysis allowed for the relative influence of multiple independent variables on the dependent variable, adjusting for any confounding effect on each other. Potential explanatory variables were entered in the model and retained in a second step if significant ($s < 0.05$). The odds ratio of each variable value was examined to determine directionality and linearity. Finally, each outcome was tabulated against the other two (time left with symptoms, time left with care options and symptoms with care options), and χ^2 tests were used to measure the association between responses with significance level of $p < 0.05$.

Results

Sample description

The demographic characteristics of the sample are displayed by country in Table 1.

From a total of 45 242 randomly selected households with a known eligible person, 9344 people (21%) agreed to participate in the study. Response was highest in Germany (29%), followed by Portugal (28%), Spain (21%), Italy (21%), England (21%), Flanders (16%) and Netherlands (16%). Overall, 1% of them were excluded because of language difficulties, 1% because of hearing problems and 1% were deemed incapable of providing informed consent.

Main reasons for refusal were lack of interest (59%), lack of time (17%) and refusal to generally take part in telephone surveys (3%). In addition, 3% ($n = 1251$) stated reasons for refusal to participate related to the topic: 497 because they had a physical disability or illness, 385 because of its sensitive nature, 223 because a relative or friend had a physical disability or illness and 146 due to a recent household bereavement/death. In 3%, reasons for refusal were not specified. In addition, 7% asked to be called back, but the interview was never concluded, 8% broke-off the interview at early stages and <1% chose to stop and withdrew their information.

The interview took 15.4 min on average to complete. The oldest person interviewed was 95 years old, 13% participants were 70 years or older and 93% were born in the country they currently lived in. Ten per cent reported having been seriously ill in the past 5 years, and 53% had cared for a close relative or friend in their last months of life (Table 2).

Descriptive analysis

Across the whole sample, 73.9% of respondents would always want to know if they had limited time left, 13.6% would only want to be told if they asked and 7.5% would not want to be informed (Table 2). More than four-fifths (81.3%) would always want to be informed about symptoms, and 89.5% would always want to be informed of their care options. The proportion always wanting to know they had limited time left varied from 67.6% in Italy to 80.7% in Flanders. The proportion wanting to know about symptoms was also lowest in Italy (76.9%) and highest in Portugal (89.6%), whereas a wish to know about care options ranged from 82.8% in the Netherlands to 94.9% in Portugal.

Associations of categorical outcomes with independent variables

With respect to age, all outcomes showed a similar distribution; in the 16–69 range there was a slight peak in the middle but little variation, while those aged 70+ were substantially less likely to want information (see Table B in supplementary material). All three outcomes were significantly associated with age ($p < 0.001$) using χ^2 tests (time left, $\chi^2 = 91.64$; symptoms, $\chi^2 = 113.92$; care options, $\chi^2 = 157.54$).

People with higher educational attainment were more likely to want to know all information, either always or if they asked, and more likely to have a view (i.e. less likely to respond 'don't know'). Education was significantly associated ($p < 0.001$) with all outcomes (time left, $\chi^2 = 139.77$; symptoms, $\chi^2 = 96.43$; care options, $\chi^2 = 101.21$).

Those living alone were less likely to want to know about symptom burden or care options. Living alone was not associated with wish to know about time left ($\chi^2 = 3.84$, $p = 0.279$) but significantly associated ($p < 0.001$) with wish to know about symptoms ($\chi^2 = 20.50$) and care ($\chi^2 = 37.30$).

Table 1. Participant characteristics by country

n	England 1351	Flanders 1269	Germany 1363	Italy 1352	Netherlands 1356	Portugal 1286	Spain 1367	All countries 9344
Age								
Mean in years (SD)	54.2 (16.3%)	52.2 (14.3%)	47.1 (15.7%)	48.7 (15.9%)	54.5 (14.6%)	50.1 (16.9%)	48.1 (16.5%)	50.7 (16.0%)
16–29	107 (8.0%)	88 (7.5%)	213 (15.8%)	177 (15.0%)	61 (4.7%)	169 (13.8%)	204 (15.4%)	1019 (11.5%)
30–39	151 (11.3%)	119 (10.2%)	197 (14.6%)	166 (14.1%)	126 (9.7%)	176 (14.4%)	213 (16.1%)	1148 (12.9%)
40–49	255 (19.1%)	261 (22.3%)	361 (26.8%)	241 (20.4%)	289 (22.3%)	231 (18.9%)	279 (21.0%)	1917 (21.6%)
50–59	258 (19.4%)	315 (27.0%)	273 (20.3%)	272 (23.1%)	313 (24.2%)	246 (20.1%)	294 (22.2%)	1971 (22.2%)
60–69	317 (23.8%)	256 (21.9%)	184 (13.7%)	209 (17.7%)	306 (23.6%)	226 (18.5%)	198 (14.9%)	1696 (19.1%)
70+	244 (18.3%)	129 (11.0%)	119 (8.8%)	115 (9.7%)	199 (15.4%)	175 (14.3%)	139 (10.5%)	1120 (12.6%)
Gender								
Female	863 (63.9%)	832 (65.6%)	790 (58.0%)	974 (72.0%)	891 (65.8%)	893 (69.4%)	935 (68.4%)	6178 (66.1%)
Living arrangements								
Living alone	325 (24.2%)	197 (15.6%)	281 (20.8%)	142 (10.5%)	294 (21.8%)	136 (10.6%)	156 (11.5%)	1531 (16.5%)
Urbanisation level								
Big city or suburbs/outskirts	500 (37.1%)	289 (22.8%)	556 (40.9%)	269 (19.9%)	363 (26.8%)	643 (50.0%)	324 (23.7%)	2944 (31.5%)
Town or small city	495 (36.7%)	224 (17.7%)	419 (30.8%)	526 (39.0%)	311 (23.0%)	368 (28.6%)	589 (43.1%)	2932 (31.4%)
Country village	287 (21.3%)	591 (46.6%)	301 (22.1%)	521 (38.6%)	578 (42.7%)	221 (17.2%)	401 (29.4%)	2900 (31.1%)
Farm or home in countryside	66 (4.9%)	165 (13.0%)	85 (6.2%)	34 (2.5%)	103 (7.6%)	54 (4.2%)	52 (3.8%)	559 (6.0%)
Marital status								
Married or with a partner	822 (61.3%)	951 (75.7%)	784 (58.1%)	860 (63.8%)	932 (69.2%)	814 (63.6%)	847 (62.2%)	6010 (64.8%)
Divorced or separated	175 (13.1%)	100 (8.0%)	152 (11.3%)	86 (6.4%)	110 (8.2%)	91 (7.1%)	100 (7.3%)	814 (8.8%)
Widowed	131 (9.8%)	96 (7.6%)	83 (6.2%)	92 (6.8%)	142 (10.5%)	109 (8.5%)	113 (8.3%)	766 (8.3%)
Single	212 (15.8%)	110 (8.8%)	330 (24.5%)	310 (23.0%)	162 (12.0%)	265 (20.7%)	301 (22.1%)	1690 (18.2%)
Religion								
Yes	778 (57.9%)	664 (52.9%)	771 (57.0%)	1094 (81.6%)	616 (45.6%)	1017 (79.6%)	959 (71.0%)	5899 (63.6%)
Health								
Very good	565 (42.0%)	490 (38.6%)	310 (22.9%)	305 (22.6%)	302 (22.3%)	170 (13.3%)	293 (21.5%)	2435 (26.1%)
Good	535 (39.8%)	583 (46.0%)	699 (51.5%)	642 (47.6%)	748 (55.3%)	488 (38.1%)	560 (41.1%)	4255 (45.7%)
Fair	191 (14.2%)	176 (13.9%)	289 (21.3%)	377 (28.0%)	254 (18.8%)	558 (43.6%)	437 (32.0%)	2282 (24.5%)
Bad	47 (3.5%)	16 (1.3%)	53 (3.9%)	21 (1.6%)	43 (3.2%)	46 (3.6%)	67 (4.9%)	293 (3.1%)
Very bad	7 (0.5%)	3 (0.2%)	5 (0.4%)	3 (0.2%)	5 (0.4%)	19 (1.5%)	7 (0.5%)	49 (0.5%)
Experience of illness, death and dying								
Diagnosed with serious illness in last 5 years	172 (12.8%)	190 (15.2%)	107 (8.0%)	113 (8.4%)	137 (10.1%)	99 (7.8%)	119 (8.8%)	937 (10.1%)
Close relative/friend seriously ill in last 5 years	849 (63.1%)	760 (60.6%)	862 (64.1%)	900 (67.4%)	969 (71.8%)	728 (57.5%)	923 (68.2%)	5991 (64.8%)
Death of close relative/friend in last 5 years	949 (70.6%)	876 (69.9%)	933 (69.4%)	928 (69.3%)	1036 (76.7%)	771 (60.9%)	1006 (74.4%)	6499 (70.3%)
Cared for close relative/friend in last months of life	679 (50.6%)	625 (49.9%)	647 (48.0%)	815 (60.8%)	702 (52.0%)	673 (53.2%)	771 (57.0%)	4912 (53.1%)

Sums may not always amount to the total sample number because of missing values on variables. Percentages may not always add up to 100 because of rounding. SD, standard deviation.

European citizens' preference to be told of poor prognosis

Table 2. Preferences by country for information on time left, likely symptoms and problems, and care options available (%)

Would want to know about	Answer options	England 1351	Flanders 1269	Germany 1363	Italy 1352	Netherlands 1356	Portugal 1286	Spain 1367	All countries 9344
Time left	Always	77.0	80.7	75.0	67.6	74.1	73.5	70.1	73.9
	If I ask	16.2	7.3	14.1	19.8	16.3	9.3	11.6	13.6
	No	5.2	6.0	5.6	7.3	6.8	10.6	11.4	7.5
	Don't know	1.6	6.1	5.3	5.4	2.8	6.6	7.0	5.0
Symptoms	Always	80.0	87.2	78.0	76.9	77.3	89.6	80.6	81.3
	If I ask	15.0	5.3	15.2	16.4	14.6	5.1	11.7	12.0
	No	3.9	4.1	4.0	4.8	6.7	2.6	5.3	4.5
	Don't know	1.1	3.4	2.8	1.9	1.4	2.7	2.4	2.2
Care options	Always	91.3	91.9	87.8	88.5	82.8	94.9	90.1	89.5
	If I ask	6.4	5.0	8.6	9.0	12.9	2.7	5.3	7.2
	No	2.0	2.1	1.8	1.9	3.3	0.8	1.8	2.0
	Don't know	0.4	1.1	1.8	0.6	1.0	1.7	2.9	1.4

Association of binary outcome with independent variables by country

The main outcome, wish to know if there was limited time left, was coded binary and tabulated against independent variables by country (Table 3).

Overall, men were more likely to wish to know, and this association was statistically significant in five countries (England, Germany, Italy, Portugal and Spain) and maintained the same direction in a sixth (Netherlands). Flanders was the exception in that women were significantly more likely to want to know about limited time left.

Widowed people were less likely to want to know if there was limited time left, overall and in five countries (although this only reached statistical significance for the Netherlands), but in Italy, the reverse was true.

Education was associated with greater probability of the outcome in all countries, reaching statistical significance in four countries (Flanders, England, Netherlands and Spain). Overall, financial hardship was associated with lower probability of the outcome, but this association was not significant in any single country.

In Italy and Portugal, participants whose main concerns were social were more likely to want to know about limited time left, but in other countries and overall, there was no association. Those who prioritised quality over extension of life had higher probability of wanting time left information in five countries (Flanders, England, Germany, Netherlands and Italy), but the reverse was true in Portugal. There was no association between this outcome and preferred place of death.

People who had been diagnosed with a serious illness in the past 5 years were more likely to want to know about limited time left in five countries (England, Germany, Netherlands, Italy and Portugal), and the other two (Flanders and Spain) showed very small differences. Severe illness in a close relative or friend was also associated with the outcome, except in Italy.

Multivariate model

Any variable associated with the outcome at a significance level of $p < 0.1$ for all countries combined, with shared directionality in a majority of countries, was carried forward into multivariate analysis. The following variables met these criteria: age, gender, marital status, education, finances, quality versus quantity of life, serious illness in self and relative with serious illness.

Age did not show a clear overall trend in bivariate analysis, but the oldest group was clearly distinct from the others, and this created significant associations. Therefore, age was analysed as categorical using dummy variables. Only the oldest age group was associated with the outcome, so the variable was recoded to binary, with 16–69 years compared to 70 years and older. Marital status was also analysed as categorical because the options were not ordered.

Wanting to know about limited time left was significantly more likely among people aged younger than 70 years (OR = 0.72), men (OR = 1.23), those with more education (OR = 1.20), those with experience of serious illness in themselves (OR = 1.20) or in others (OR = 1.12), and those who prioritised quality over quantity of life (OR = 1.13) (Table 4). Therefore, age had the greatest effect. The difference between countries was statistically significant ($p < 0.001$).

Associations between outcomes

There was a high degree of association between the three outcomes (χ^2 over 1000 in all three tests, $p < 0.001$). Participants who wanted information in any area were more likely to want information in the other two (Table 5). However, many people wanted selective information. Of those who did not want to be told if they had limited time left, less than half (41.8%) wished to be told their probable symptoms, but two-thirds (67.1%) wanted to know their care options. The question regarding time left had the highest prevalence of 'don't know' responses in the entire questionnaire. Participants who did not know whether they would want to be informed if they had limited time left were likely to say they would always want to know their symptoms (58.2%) and care options (81.3%).

Discussion

Our data have revealed an international majority public preference to always be informed in the scenario of having a serious illness with less than a year to live. However, the preference to know about remaining time left (overall 73.9%) was smaller than both desire for knowledge about possible symptoms (81.3%) and for care options (89.5%). Although a strong correlation was found between the three outcomes, even among those who would not want to know if they had limited time left, the majority (67.1%) would always want to know the care options available to them.

Table 3. Crude percentages of wish to know about time left by covariates and countries

	England	Flanders	Germany	Italy	Netherlands	Portugal	Spain	All countries
<i>Sociodemographic variables</i>								
Age bands (years)		*			*	*		***
16–29	76.6	81.8	69.8	67.8	67.2	79.9	71.1	73.0
30–39	82.1	79.0	74.4	69.3	70.4	74.9	74.7	74.8
40–49	80.8	85.4	76.6	65.2	77.2	75.2	70.6	76.0
50–59	74.0	81.3	78.7	67.2	78.5	71.0	68.2	74.4
60–69	78.1	78.5	76.8	64.4	74.5	73.0	71.1	74.2
70+	71.7	71.9	71.4	64.4	65.7	63.4	63.0	67.5
Gender	*	*	*	*		*	*	***
Male	80.6	77.1	78.0	71.7	75.2	77.8	74.0	76.5
Female	74.9	82.7	72.9	66.0	73.6	71.6	68.2	72.6
Marital status							*	**
Married or with civil partner	78.7	81.4	76.9	68.2	74.5	72.5	69.0	74.5
Divorced or separated	75.4	82.0	72.9	65.9	79.1	79.1	77.0	75.9
Widowed	68.7	78.1	67.1	72.8	68.1	67.9	62.0	69.0
Single	76.8	76.4	74.1	64.5	74.1	76.6	73.8	73.2
Highest level of education	*	**			*		**	***
Up to 16	68.8	65.4	33.3	40.0	58.3	66.7	60.2	64.3
16–18	78.2	78.8	74.5	68.0	72.3	73.5	68.5	72.8
18+	78.0	84.0	76.6	65.8	78.1	73.9	75.9	77.0
Financial hardship								**
Difficult or very difficult on present income	74.7	79.4	71.8	68.3	70.7	76.3	66.8	72.1
Coping on present income	74.6	78.8	77.5	68.9	73.2	72.3	68.8	73.2
Living comfortably on present income	80.2	82.3	73.4	65.1	75.1	73.9	74.0	75.6
<i>Value/preference-based variables</i>								
Most concerning problem				*		*		
Social focussed concern	76.1	81.4	75.7	71.2	74.3	79.1	71.4	74.9
Physical/psychological concern	77.8	81.0	74.8	65.5	74.1	72.6	70.2	74.1
Quantity/quality of life								**
Extend life	63.3	74.6	74.5	63.9	71.7	78.2	69.1	71.8
Both are equally important	75.2	78.1	71.7	66.0	68.9	74.1	71.4	71.7
To improve the quality of time left	78.0	82.0	77.4	68.7	76.0	73.5	70.4	75.2
Preferred place of death								
Own home/home of relative or friend	78.7	79.2	76.4	67.5	74.5	73.7	70.6	74.2
Hospice or palliative care unit	76.9	84.4	75.1	69.2	72.5	76.1	73.8	75.5
Hospital – but not a palliative care unit	73.8	83.5	64.4	67.1	72.1	67.3	70.0	72.4
Nursing home or residential home	65.4	87.5	75.0	63.6	81.8	73.1	51.4	72.9
Somewhere else	56.5	66.7	78.0	66.7	88.9	78.8	66.7	71.4
<i>Experiential variables</i>								
Self diagnosed with serious illness in last 5 years					*			**
Yes	83.7	80.4	78.3	73.2	77.4	77.6	69.5	77.8
No	75.9	80.8	74.8	67.2	73.8	73.1	70.4	73.6
Close relative/friend diagnosed with serious illness in the last 5 years		***						*
Yes	77.8	84.2	76.2	66.4	74.1	75.0	71.6	74.8
No	75.2	75.5	72.8	71.0	74.3	71.4	67.3	72.5

Sums may not always amount to the totals because of missing values on variables.

* $p < 0.05$.

** $p < 0.01$.

*** $p < 0.001$.

Table 4. Multivariate logit model of wish to know about time left

Covariate	Reference group	OR	95% CI	Z	s
Age	16–69 years	0.72	0.62–0.83	–4.44	<0.001
Gender	Female	1.23	1.10–1.37	3.77	<0.001
Education	16–18 years	1.20	1.09–1.32	3.83	<0.001
Serious illness	No	1.20	1.01–1.43	2.05	0.040
Relative's illness	No	1.12	1.01–1.24	2.15	0.032
Quantity or quality of life	Quantity	1.13	1.03–1.24	2.69	0.007

These data provide important evidence that clinicians should anticipate patients wanting full knowledge on their prognosis, disease presentation and care (although a minority varying by country do not wish to know unless they ask, or at all, with a quarter overall not wanting information on prognosis). The analysis also found that people

younger than 70 years, men, those with experience of illness and with higher educational attainment were more likely to want to know if they had limited time left.

It is notable that the three countries with the lowest percentage of respondents who would wish to know about time left were Italy, Spain and Portugal. It has been suggested that in traditional Mediterranean Catholic countries, patients are more likely to reject autonomy and that the medical establishment may have a greater paternalism [20].

These findings now demand evidence of appropriate and effective communication skills to enable clinicians to elicit preferences and of information-giving. A systematic review of clinician–patient communication [21] found little definitive evidence to support one communication method over any other. Clinicians should be adequately equipped from

European citizens' preference to be told of poor prognosis

Table 5. Associations between information preference items

Preference to know	N	Symptoms %				Care %				
		Always	If I ask	No	D/K	Always	If I ask	No	D/K	
Time left	Always	6880	92.0	5.3	1.7	1.1	94.2	4.1	1.0	0.7
	If I ask	1264	53.5	41.6	3.3	1.6	79.4	18.3	1.0	1.3
	No	699	41.8	20.5	31.9	5.9	67.1	15.7	13.3	3.9
	D/K	460	58.2	18.8	7.4	15.5	81.3	9.6	1.5	7.6
Care	Always	8338	86.5	9.4	2.6	1.6	—	—	—	—
	If I ask	666	38.1	46.1	12.0	3.8	—	—	—	—
	No	181	30.9	7.2	56.4	5.5	—	—	—	—
	D/K	125	39.2	12.8	15.2	32.8	—	—	—	—

D/K = don't know.

medical and nursing training curricula to manage information disclosure. However, evidence suggests that although communication training is effective in terms of skill acquisition, this is less well applied in practice [22].

The skill needed in eliciting preferences and sharing information also needs to extend to those who may not be clear on whether they would want information to be shared with them. One in 20 people did not know whether they would want to know about time left, and this was less common for other questions. That people were unable to make a hypothetical decision does not necessarily translate to a real situation of having a serious illness such as cancer. However, it is interesting that in the multivariate analysis, personal experience of a serious illness increased the likelihood of wanting to know about time left.

Age was a very important covariate in our place of death study [10]. Three effects are potentially related to each other with respect to birth cohort effect, health status and covariates. Older people are more likely to have health problems, to be retired and widowed. It is difficult to disentangle these effects.

These findings derive from a robust cross-national comparison. We used standard methodologies and asked identical questions across countries (reducing differential biases). The results provide not only country-specific factors and local preferences (important since most policies are made at national level) but also cross-national data to inform international policy making.

However, the study has also some limitations. Our response rate is typical of the declining rates of random digit dialling surveys [23], although in some countries, our response rate is higher than the rates in the 2002 Picker Institute Europe survey of public views on health care responsiveness [24]. Our sample presents well-known selection biases excluding those living in households without a fixed telephone (29% of households in the EU-27) [24] and over-representing women and older people because of selective non-response. The bias towards women may have reduced estimates, as men were more likely to want information on time left. Notwithstanding, these limitations have little effect on the main aim of cross-national comparisons.

For some of the healthier participants and as identified in the pilot stages of the questionnaire [14], it might have been difficult to imagine the actual circumstances surrounding the dying process and preceding months. However we aimed to minimise this through the use of a well-conceived hypothetical scenario. Also, this might have been easier for people with a previous experience of caring for someone close to them in the last months of life (53% of our sample). Experiences of a diagnosis

of serious illness in self or in another were associated with the outcome of wanting to know about time left, suggesting that those who made a hypothetical choice in the absence of experience may be likely to choose information in the presence of serious illness. Data from Australian patients with metastatic disease reported a slightly lower preference than that reported in our survey (65% vs 73.9%) to know about 1-year survival [25]. Data from Brazilian patients with advanced cancer found that the vast majority (92%) believed that patients should know their poor prognosis, although interestingly, only around one-third had been informed themselves [26]. A further study from Korea of advanced cancer patients found a majority preference for prognosis disclosure (78.6%) [27]. However, among cancer outpatients in Japan, a study reported only 50.4% of the sample reported preferring to know about time left [28]. Therefore, it seems that the hypothetical/real scenario may be less clearly related to preference than culture and context. We also recognise that although there may be a preference to know about poor prognosis, the evidence suggests that accurate prognostication is difficult [29].

The country models helped understand within-country diversity. The models confirmed the influence of four factors in more than one country (age, gender, education and most concerning problem) and added 11 country-specific factors to which national policies and clinical practice should be responsive.

Replicating the survey with patients at different stages of illness would help determine if and how hypothetical and 'real' preferences differ. Policy should now reflect the majority preference for full information on time left, likely symptoms and care options, as well as putting adequate care options in place for patients with advanced disease.

Supporting information

Supporting information may be found in the online version of this article.

Acknowledgements

We are most grateful to all the survey participants. We thank the European Commission for the financial support needed to undertake this study, BMG Research and ZEM University of Bonn for assistance in survey administration and data collection, and Gao Wei and Joana Cadima for statistical advice. We also thank our colleagues from PRISMA including the scientific committee who contributed to discussions and scientific review of the survey, namely, Hamid Benalia, Emma Bennett, Lucy Bradley, Noël Derycke, Martine de Vlioger, Let Dillen, Natalie Evans, Michael Ehteld, Nancy Gikaara,

Stein Kaasa, Johan Menten, Bregje Onwuteaka-Philipsen, Ana Barros Pinto, Robert Pool, Richard A. Powell, Miel Ribbe, Katrin Sigurdardottir, Bart Van den Eynden, Paul Vanden Berghen and Trudie van Iersel. We thank Susana Bento, Carolina Comabella, Filomena Ferreira, Grethe Iversen, Carmen López-Dóriga, Constanze Rémi, Christian Schulz and Wessex Translations for their work translating and back-translating the questionnaires. The invaluable work of Ron Irwin, Sian Best and Mike Gover at King's College London is also highly appreciated. This work was supported by the European Commission's Seventh Framework Programme (contract number: Health-F2-2008-201655). PRISMA was funded by the European Commission's Seventh Framework Programme with the overall aim to co-ordinate high-quality international research into end-of-life cancer care. PRISMA aims to provide evidence and guidance on best practice to ensure that research can measure and improve outcomes

for patients and families. PRISMA activities aim to reflect the preferences and cultural diversities of citizens, the clinical priorities of clinicians, and appropriately measure multidimensional outcomes across settings where EoLC is delivered. Principal Investigator: Richard Harding. Scientific Director: Irene J Higginson. This article reflects only the author's views and the European Commission is not liable for any use that may be made of the information contained therein. The funder had no role in the study design, in the collection, analysis and interpretation of data, in the writing of this manuscript, or in the decision to submit this manuscript for publication.

Conflict of interest

The authors have declared that there is no conflict of interest.

References

- Higginson IJ, Costantini M. Dying with cancer, living well with advanced cancer. *Eur J Cancer* 2008;**44**(10):1414–1424.
- Wenrich MD *et al.* Communicating with dying patients within the spectrum of medical care from terminal diagnosis to death. *Arch Intern Med* 2001;**161**(6):868–874.
- Clayton JM, Butow PN, Tattersall MHN. When and how to initiate discussion about prognosis and end-of-life issues with terminally ill patients. *J Pain Symptom Manage* 2005;**30**(2):132–144.
- Street Jr RL. Information-giving in medical consultations: the influence of patients' communicative styles and personal characteristics. *Soc Sci Med*, 1991;**32**(5):541–548.
- Fitch MI. How much should I say to whom? *J Palliat Care* 1994;**10**(3):90–100.
- Schulman-Green D *et al.* Nurses' communication of prognosis and implications for hospice referral: a study of nurses caring for terminally ill hospitalized patients. *Am J Crit Care* 2005;**14**(1):64–70.
- Mackillop WJ *et al.* Cancer patients' perceptions of their disease and its treatment. *Br J Cancer* 1988;**58**(3):355–358.
- Gaston CM, Mitchell G. Information giving and decision-making in patients with advanced cancer: a systematic review. *Soc Sci Med* 2005;**61**(10):2252–2264.
- Mauri E *et al.* An exploratory study on the Italian patients' preferences regarding how they would like to be told about their cancer. *Support Care Cancer* 2009;**17**(12):1523–1530.
- Gomes B *et al.* Preferences for Place of Death if Faced with Advanced Cancer: A Population Survey in England, Flanders, Germany, Italy, the Netherlands, Portugal and Spain. *Ann Oncol* 2012.
- Harding R, Higginson IJ. PRISMA: a pan-European co-ordinating action to advance the science in end-of-life cancer care. *Eur J Cancer* 2010;**46**(9):1493–1501.
- Jowell R. *Measuring Attitudes Cross-nationally: Lessons from the European Social Survey*: Sage: London 2007.
- Cull A *et al.* EORTC quality of life group translation procedure, *EORTC Quality of Life Group Publication*: Brussels 2002.
- Daveson BA, *et al.* Constructing understandings of end-of-life care in Europe: a qualitative study involving cognitive interviewing with implications for cross-national surveys. *J Palliat Med* 2011;**14**(3):343–349.
- Hall S *et al.* *Palliative Care for Older People: Better Practices*. W.H.O.R.O.f. Europe, Editor: Copenhagen 2011.
- Cohen J *et al.* Which patients with cancer die at home? A study of six European countries using death certificate data. *J Clin Oncol* 2010;**28**(13):2267–2273.
- Inglehart R, Welzel C. *Changing Mass Priorities: The Link Between Modernization and Democracy*. Perspectives on Politics 2010;**8**(2):551–567.
- Busse R *et al.* Tackling chronic disease in Europe: strategies, interventions and challenges. in *Observatory Study Series*, <http://www.cureacasa.it/files/bio2/10.pdf>, WHO Regional Office for Europe: Copenhagen 2010.
- Higginson I. *Priorities and Preferences for End-of-life Care*: London 2003.
- Toscani F, Farsides C. Deception, catholicism, and hope: understanding problems in the communication of unfavorable prognoses in traditionally-catholic countries. *Am J Bioeth* 2006;**6**(1):W6–18.
- Rodin G *et al.* Clinician-patient communication: a systematic review. *Support Care Cancer* 2009;**17**(6):627–644.
- Gysels M, Richardson A, Higginson IJ. Communication training for health professionals who care for patients with cancer: a systematic review of effectiveness. *Support Care Cancer* 2004;**12**(10):692–700.
- Kempf AM, Remington PL. New challenges for telephone survey research in the twenty-first century. *Annu Rev Public Health* 2007;**28**:113–126.
- Communication., D.-G.f., E-communications household survey, in *Special Eurobarometer 293*, http://ec.europa.eu/public_opinion/archives/ebs/ebs_293_full_en.pdf, Editor 2008.
- Hagerty RG *et al.* Cancer patient preferences for communication of prognosis in the metastatic setting. *J Clin Oncol* 2004;**22**(9):1721–1730.
- Fumis RR, De Camargo B, Del Giglio A. Physician, patient and family attitudes regarding information on prognosis: a Brazilian survey. *Ann Oncol* 2012;**23**(1):205–211.
- Yun YH *et al.* Experiences and attitudes of patients with terminal cancer and their family caregivers toward the disclosure of terminal illness. *J Clin Oncol* 2010;**28**(11):1950–1957.
- Fujimori M *et al.* Preferences of cancer patients regarding the disclosure of bad news. *Psycho-Oncology* 2007;**16**(6):573–581.
- Moss AH *et al.* Prognostic significance of the "surprise" question in cancer patients. *J Palliat Med* 2010;**13**(7):837–840.