

PROTOZOA

Taxonomy - classification


- A. **Amebas** - pseudopodia – moving and feeding. The **important pathogens** belonging to this group are ***Entamoeba histolytica*, *Naegleria fowleri*** and ***Acanthameba***.
- B. **Flagellates** – flagella – moving, cytostome - feeding. ***Trichomonas vaginalis*, *Giardia lamblia*, *Trypanosoma*** species and ***Leishmania*** species belong to this group.
- C. **Ciliates** - move by beating of cilia synchronously and have a more complex digestive system (***Balantidium coli*** – rare human GIT infection).
- D. **Sporozoa** (apicomplexa) - are obligate intracellular parasites that have a complex life cycle with more than one host and generally have a non motile adult form. Pathogens belonging to this group are: ***Cryptosporidium*, *Plasmodium* species and *Toxoplasma gondii***.

Amebas – infections, diagnosis, treatment

Amebas - pseudopodia – moving and feeding. The **important pathogens** belonging to this group are *Entamoeba histolytica*, *Naegleria fowleri* and *Acanthameba*.

- *Entamoeba histolytica* – Amebic dysentery or liver abscess, source – cysts in contaminated water or food form **trophozoites in the small intestine**, in colone feed on bacteria and may invade epithelium (ulceration), **can spread to the liver (abscesses)**
- **Dg – fecals samples** – motile **trophozoites**, **cysts** or specific DNA (**PCR**) in fecal samples, **liver abscess** – serum to detect **IgG specific antibodies**
- **Therapy – metronidazol**

Amebas – cycle



trophozoite



cyst

<https://www.cdc.gov/dpdx/amebiasis/index.html>

Other ameboid


Naegleria fowleri (commonly referred to as the “**brain-eating amoeba**”), is a **free-living** microscopic ameba, (single-celled living organism). It can cause a **rare and devastating infection** of the brain called **primary amebic meningoencephalitis (PAM)**. The ameba is commonly found in warm **freshwater** (e.g. lakes, rivers, and hot springs) and soil. **Source:** contaminated water **enters the body through the nose** and **travels to the brain** where it causes **PAM**, which is **usually fatal**. Infection typically occurs when people go **swimming** in warm **freshwater** places, like **lakes** and **rivers**. In very rare instances, *Naegleria* infections may also occur when contaminated water from other sources (such as inadequately chlorinated **swimming pool** water or heated and **contaminated tap water**) enters the nose. You **cannot** get infected from **swallowing** water contaminated with *Naegleria*.

Diagnosis – direct visualisation or PCR from cerebrospinal fluid

The treatment of choice for primary amoebic meningoencephalitis (PAM), or infection with brain-eating amoeba (*Naegleria fowleri*) is the **antifungal amphotericin B**. Some survivors in North America were treated with a combination of drugs that included amphotericin B, rifampin, fluconazole and a drug called miltefosine.

DIAGNOSIS – see also <https://www.cdc.gov/parasites/naegleria/diagnosis-hcp.html>

Naegleria fowleri



Computer-generated representation of Naegleria fowleri in its ameboid trophozoite stage, in its flagellated stage, and in its cyst stage. (CDC, Atlanta)

Acanthamoeba

Acanthamoeba is a microscopic, **free-living ameba**, that can cause **rare, but severe infections of the eye, skin, and central nervous system**. The ameba is found worldwide in the environment in water and soil. The ameba can be spread to the eyes through **contact lens use, cuts, or skin wounds** or by being inhaled into the lungs. Most people will be exposed to *Acanthamoeba* during their lifetime, but very few will become sick from this exposure.

***Acanthamoeba* keratitis** – An infection of the eye that typically occurs in healthy persons and can result in **permanent visual impairment or blindness**.

Granulomatous Amebic Encephalitis (GAE) – A serious infection of the brain and spinal cord that typically occurs in **persons with a compromised immune system**.

Disseminated infection – A widespread infection that can affect the skin, sinuses, lungs, and **other organs** independently or in combination. It is also more common in persons with a **compromised immune system**.

Diagnosis – clinically, scraping of the eye – **direct detection** by confocal **microscopy** or **PCR**

Treatment – **antiseptics** (e.g. chlorhexidine) and **antibiotics** (e.g. neomycine) (more details see also: <https://www.aao.org/education/current-insight/acanthamoeba-keratitis-treatment>)

Acanthamoeba keratitis



CASE REPORT: <https://www.aao.org/education/clinical-video/acanthamoeba-revenge>

Acanthamoeba keratitis (AK) is a **corneal infection characterized by an indolent course and challenging management**. Known risk factors include **contact lens (CL) wear, freshwater exposure, and trauma**. Dr. Jaime Martinez and Pietro Carletti describe the case of a 55-year-old woman with a history of CL use who presented with a **nonresolving corneal ulcer**. Three weeks after onset, cultures demonstrated no growth, and AK was diagnosed with a corneal biopsy. Her condition improved initially with long-term maximal topical therapy, but then recurred and was refractory to aggressive treatment, including oral miltefosine and Rose Bengal photodynamic antimicrobial therapy. She progressed to develop scleral involvement, elevated intraocular pressures, and severe pain, and thus underwent therapeutic **corneoscleral transplantation** and open-sky cataract extraction. At 2 months after surgery, there were no signs of infection recurrence.

Flagellates – infections, diagnosis, treatment


Giardia lamblia

- Two forms: cyst and motile trophozoite
- gastric acid stimulates excystation
- release **trophozoites in duodenum** that **adhere to small intestine wall**



Interferes fat absorption
Steatorrhea

See also: <https://www.youtube.com/watch?v=Rjw6kWiRX8Y>
<https://www centralsan.org/post/giardia-lamblia>



Giardia lamblia


- **Fat packed stools** gives giardia it's best known symptom: **Foul smelling diarrhea!**
- Giardia **does not invade the intestinal wall!** There is **no blood in the stool!**
- **Epidemiology:** worldwide, sylvatic, streams, lakes
- **Source - water, uncooked vegetables or fruits, fecal-oral transmission**
- **cyst stage is resistant to chlorine** concentrations used in most water-treatment facilities
- asymptomatic carriage (50%) or symptomatic disease, ranging **from mild diarrhea to severe malabsorption syndrome**
- **Long incubation 1-4 weeks, sudden onset**
- **Usually spontaneous recovery** after 10-14d, although more chronic disease with multiple relapses - particularly pp with Ig A deficiency or intestinal diverticula.

Giardia lamblia

- **Diagnosis:**
 1. Examination of **stool** specimens for **cysts/ trophozoites or specific DNA (PCR)**
 2. You should never accept single negative
 3. 1 specimen/day for **3 days should be examined**
- If remain negative and giardiasis is highly suspected: duodenal aspiration sample is suggested
- Immunologic tests for fecal antigen: highly sensitive and specific
- **Treatment: metronidazole**

Trichomonas vaginalis

- Flagellate, 4 **short flagella**, **undulating membrane**, Trophozoite form only
- **Disease: STD**
- men: urethra + prostate
- women: urethra + vagina
- Clinical:
- **women**: asymptomatic or scant, watery vaginal discharge;
- **vaginitis** - extensive inflammation (strawberry vagina)
- malodorous, yellow green **vaginal discharge**
- erosion of epithelial lining – **itching, burning and painful urination**
- **men**: mainly asymptomatic carriers
- occasionally: **urethritis**, prostatitis, other urinary problems




Trichomonas vaginalis

- **Epidemiology:** worldwide; developed countries: 5-20% women; 2-10% men;
- **Transmission:**
 - **Sexual intercourse**
 - Fomites (eg.clothing)
 - Birth canal infection
- **Laboratory Diagnostics:**
 - **microscopic examination** of urethral/ vaginal discharge – unstained or **stained with Giemsa** – motile trophozoites
 - Improved by **culture**
 - **urine - PCR**
- **Treatment: metronidazole** – patient + sexual partner
- See also: <https://www.youtube.com/watch?v=rin2dXF3Oac>


Leishmania (*L. donovani*, *L. tropica*, *L. mexicana*)

Leishmaniasis is a parasitic disease that is found in parts of the **tropics, subtropics**, and **southern Europe**. Leishmaniasis is caused by infection with *Leishmania* parasites, which are **spread** by the **bite** of **phlebotomine sand flies**. There are several different forms of leishmaniasis in people. The most common forms are **cutaneous leishmaniasis**, which causes skin sores, and **visceral leishmaniasis**, which affects several internal organs (usually spleen, liver, and bone marrow).



The sand flies that transmit the parasite are only about **one third the size of typical mosquitoes or even smaller**. On the left, an example of a vector sand fly (*Phlebotomus papatasi*) is shown; its blood meal is visible in its distended transparent abdomen. On the right, ***Leishmania* promastigotes from a culture are shown**. The flagellated promastigote stage of the parasite is found in sand flies and in cultures. (CDC, Atlanta)

Leishmania life cycle



Leishmaniasis is transmitted by the bite of infected **female phlebotomine sandflies**. The sandflies inject the infective stage (i.e., promastigotes) from their proboscis during blood meals. The **number 1**. **Promastigotes** that reach the puncture wound are **phagocytized by macrophages (number 2)** and other types of mononuclear phagocytic cells. Promastigotes transform in these cells into the tissue stage of the parasite (i.e., **amastigotes (number 3)**), which multiply by simple division and proceed to **infect other mononuclear phagocytic cells (number 4)**. Parasite, host, and other factors affect whether the infection becomes symptomatic and whether **cutaneous** or **visceral** leishmaniasis results. **Sandflies become infected by ingesting infected cells during blood meals (number 5, 6)**. In sandflies, amastigotes transform into promastigotes, develop in the gut (number 7) and migrate to the proboscis (number 8).

Life cycle see also: <https://www.youtube.com/watch?v=rpxv4dhbYgk>

See also: <https://www.infond.org/practical-material/parasitic-diseases-lectures-8-visceral-leishmaniasis>

Infections

The most common form is **cutaneous leishmaniasis**, which causes **skin sores**. The sores typically develop within **a few weeks or months of the sand fly bite**. The sores can change in size and appearance over time. The sores may start out as **papules** (bumps) or **nodules** (lumps) and may end up as **ulcers** (like a volcano, with a raised edge and central crater); skin ulcers might be covered by **scab** or **crust**. The sores usually are painless but can be painful. Some people have **swollen glands near the sores**.

The other main form is **visceral leishmaniasis**, which **affects** several **internal organs** (usually **spleen, liver, and bone marrow**) and can be life threatening. The illness typically **develops within months** (sometimes as long as years) of the sand fly bite. Affected people usually have fever, weight loss, **enlargement (swelling) of the spleen and liver**, and **low blood counts** - anemia, leukopenia and thrombocytopenia.



Mucosal leishmaniasis is an example of one of the **less common forms of leishmaniasis**. This form can be a sequela of infection with some of the species (types) of the parasite that cause cutaneous leishmaniasis in parts of Latin America: certain types of the **parasite** might **spread** from the skin and **cause sores** in the **mucous membranes** of the nose (most common location), mouth, or throat. The best way to prevent mucosal leishmaniasis is to ensure adequate treatment of the original cutaneous (skin) infection.



Marked splenomegaly (enlargement/swelling of the spleen) in a patient in lowland Nepal who has **visceral leishmaniasis**. (Credit: C. Bern, CDC)



Ulcerative skin lesion, with a raised outer border, on a Guatemalan patient who has **cutaneous leishmaniasis**. (Credit: B. Arana, MERTU, Guatemala)




Mucosal leishmaniasis Lesions may multiply and increase in size, which can contribute to **severe deformity**. Because it causes major disability, those infected may be humiliated and suffer from being **ostracized by their society**. Respiratory tract mucosal invasion may also occur, causing numerous respiratory problems, and can result in malnutrition and pneumonia. Secondary infection is responsible for most deaths

(<https://web.stanford.edu/class/humbio103/ParaSites2006/Leishmaniasis/Mucocutaneous.htm>).

Diagnosis and therapy

Various laboratory methods can be used to diagnose leishmaniasis—to detect the parasite as well as to identify the *Leishmania* species (type). Some of the methods are available only in reference laboratories. In the United States, CDC staff can assist with the testing for leishmaniasis. **Tissue specimens**—such as from **skin sores** (for cutaneous leishmaniasis) or from **bone marrow** (for visceral leishmaniasis)—can be examined for the parasite under a **microscope**, in special **cultures**, and **in other ways (e.g. PCR)**. Blood tests that **detect antibody** (an immune response) to the parasite can be helpful for cases of **visceral leishmaniasis**; tests to look for the parasite itself usually also are done.



Light-microscopic examination of a stained bone marrow specimen from a patient with visceral leishmaniasis—showing a macrophage (a special type of white blood cell) containing multiple *Leishmania amastigotes* (the tissue stage of the parasite). Note that each amastigote has a **nucleus** (red arrow) and a rod-shaped **kinetoplast** (black arrow). Visualization of the kinetoplast is important for diagnostic purposes, to be confident the patient has leishmaniasis. (CDC Atlanta)

Therapy – sodium stibogluconate with pentamidine and amphotericin B


Sporozoa (apicomplexa) – GIT infections, diagnosis, treatment

Cryptosporidium

- Intestinal apicomplexa
- **Ingested as round oocyst** with 4 motile sporozoites
- Life cycle occurs **within intestinal epithelial cells** and produce more infectious oocysts



Causes **diarrhea** and **abdominal pain**



Cryptosporidium

- In **immunocompetent patient** – **self-limiting**
- In **immunocompromised** – **severe diarrhea** that is **life-threatening** (3-17L of stools per day!!)
- **Epidemiology:**
 - worldwide
 - **Fecal-oral** usually by **waterborne transmission**
 - Resistant to usual water-purification procedures (chlorination and ozone); zoonotic spread;
 - Risk: veterinarians, animal handlers, homosexuals; municipal swimming pools and day-care centers

Cryptosporidium

- **Diagnosis:**

- **Acid-fast** oocysts in stool sample
- **PCR** – DNA isolates from **stool sample**
- **Minimum 3 specimens** should be examined

- **Therapy:**

- Main goal is restoration of immune function
- Fluid replacement
- Nitazoxanide or paramomycin can be tried


Toxoplasma gondii

- **Coccidian** parasite, phylum ***Sporozoa***.
- **Intracellular parasite** in animals (including birds) and humans
- Reservoir is household **cat** (contact with cat feces)
- Many other animal sources. Humans into contact with:
 - 1) **Improperly cooked meat** of animal origin that are intermediate hosts.
 - 2) **Ingestion of infective oocysts**

Toxoplasma gondii

- **Clinical:**
- TRANSPLACENTAL infection (vertical), also transfusions and IV drug users.
- Most inf'n asymp. Problem when it relocates from blood to tissue (becoming an intracellular parasite)
- Lung, heart, lymphoid organs and CNS.

Toxoplasma gondii



Toxoplasma pathogenesis: <https://www.youtube.com/watch?v=StStwxgvuJM>

Toxoplasma gondii

- **Signs and symptoms:**
 - Acute- rash, fever, chills.
 - Chronic- hepatitis, lymphadenitis, myocarditis and chorioretinitis -> blindness.
- **1st Tri= Spontaneous abortion or still birth**
- **After 1st: Asymp but then: Epilepsy, encephalitis, microcephaly, hydrocephalus, chorioretinitis**

Toxoplasma gondii

- **Diagnosis:**
- **Serology for acute infections** (serially collected bloods with increasing Ab titres).
- Immunocompetent patients and pregnant women: test **IgM and IgG Abs to *T.gondii***. Confirmation - at a Toxoplasma reference lab.
- **Imaging methods** (e.g.MRI) to detect **toxopnecephalitis**
- **PCR from affected tissue**

Toxoplasma gondii

- **Treatment:**
- Most mononucleosis-like infections resolve spontaneously and require no therapy.
- **Always treat disseminated or CNS toxo.**
- HIV setting means high dose of **pyrimethamine** and **sulphadiazine** and then low dose forever.
- **co-trimoxazole/trimethaprim sulpha**

Trypanosomes

- Two distinct disease forms.

1. ***T. brucei gambiense*** and ***T. brucei rhodiense***

- Tsetse fly vector >> African sleeping sickness

2. ***T. cruzi***

- Reduviids >> Chagas disease (American)

Trypanosoma brucei gambiense

- Trypomastigote is infective stage, present in salivary glands of transmitting **Tsetse flies**.
- Has free flagellum and undulating membrane.
- Moves into the organism through bite wound of the fly, into CNS-reproduce in blood, lymph and CSF by binary or longitudinal fission.
- Another Tsetse infected when bites human
- Epimastigote develops into Trypo after 4-6wks

Trypanosoma brucei gambiense

Clinical:

- **Ulcer at bite site.** Chronic course, usually fatal. Lymph nodes invaded with swelling of post. cervical LNs typical in Gambian.
- **Mental retardation, meningoencephalitis.** Then convulsions, hemiplegia and incontinence.
- **Death due to CNS damage** or infections (malaria/pneum)

Trypanosoma brucei gambiense

Lab diagnosis:

- **Thick and thin films** in concentrated anti-coagulated blood preparations and LN and CSF aspirations. Centrifugation and anion-exchange chromatography.
- **ELISA** or **immunofluorescence** but costly and reagents can't be bought

Trypanosoma brucei gambiense

Treatment:

- Suramin treats acute blood and lymphatic stages of the dz. Suramin and alternative Pentamidine don't cross BBB. So Melarsoprol is used in CNS involvement.
- DFMO (Difluoromethylornithine) is a cytostatic used for acute and late (CNS) stages.
- Use of traps and insecticide with the drugs most effective.

Trypanosoma brucei rhodesiense

- Similar life cycle to T.b.gambiense.
- Incubation shorter.
- Cattle and sheep are hosts and wild game animals reservoir hosts. Makes it more difficult to control!
- Acute illness (fever, rigors and myalgia)
- 9-12 months dead

Trypanosoma brucei rhodesiense

Lab diagnosis:

- Blood and CSF like T.b.gambiense

Treatment:

- The same again!

Trypanosoma cruzi

- Additional Amastigote- no flagellum and no undulating membrane.
- Infective trypomastigote found in faeces of Reduviid 'kissing bug' enters wound. Mouth usually site of bite!
- Trypomastigote has tropism to cardiac muscle, liver and brain -> Loses flagellum and undulating memb -> Into Amastigote

Trypanosoma cruzi

- Chagoma at bite site and oedema and rash around eyes and face as Romana sign.
- Lacking agents against Chagas'
- Benzidazole and Nifurtimox for acute but not so much for chronic.
- Major side effects.
- DDT for everyone means less malaria and Chagas'.

Chagoma



Sporozoa (apicomplexa) – blood/tissue infections, diagnosis, treatment


Plasmodium

- **2 hosts:** *Anopheles* mosquito (sexual) & humans/animals (asexual)
- Infection of **300-500 million people worldwide/ year**
- **1-3 million deaths/year** (85% Africa);
- **Asexual life cycle in human cells:** *Plasmodium* sporozoites (thin, motile, spindle-shaped) in *Anopheles* saliva injected into the bloodstream, rapidly migrate to the liver to form schizonts (cyst-like structure) releasing thousands of merozoites into the bloodstream, which invade red blood cells (RBCs) using hemoglobin as nutrients. Infected RBCs rupture releasing again merozoites (through trophozoites, schizonts). If large number of RBCs rupture at the same time **paroxysm** (sudden onset) can result from the massive release of toxic substances.

See also the review article: <https://www.nature.com/articles/nrdp201750>

See also video: <https://en.wikipedia.org/wiki/Malaria>

The Life Cycle of Malaria



- 1 To start the cycle, an infected female *Anopheles* mosquito injects sporozoites into the skin while feeding.

An infected mosquito starts the cycle



SPOROZOITES





- 2 Sporozoites enter the blood stream and are carried to the liver, where they infect liver cells.


- 3 Within liver cells, the parasites develop into schizonts.

- 4 The schizonts rupture, releasing thousands of individual merozoites into the bloodstream.


HUMAN LIVER STAGES
(About 2 weeks)



Another mosquito becomes infected, continuing the cycle



- 6 Some parasites change into male and female forms called gametocytes.




- 7 When the mosquito feeds, gametocytes are ingested into its stomach.
- 8 The gametocytes emerge from the infected blood cells, becoming gametes.
- 9 The male gamete fuses with the female gamete producing a zygote.
- 10 The zygotes elongate into ookinets which move through the stomach wall.
- 11 The ookinets develop into oocysts.
- 12 The oocysts grow and rupture, releasing sporozoites.
- 13 The sporozoites migrate to the salivary glands, ready to be injected and renew the cycle.

See also:
<https://www.youtube.com/watch?v=QGxn6MEmlnY>

Malaria is endemic

- The mosquitoes that spread malaria are found in Africa, Central and South America, parts of the Caribbean, Asia, Eastern Europe, and the South Pacific



This map shows an approximation of the parts of the world where malaria transmission occurs.

<https://www.cdc.gov/malaria/about/distribution.html>

Sporozoa (apicomplexa) – blood/tissue infections, diagnosis, treatment

Plasmodium

- *P. vivax* & *P. ovale* establish dormant stage: **hypnozoites**
- attach/ enter RBCs: **erythrocytic cycle**
- 3 phases of **tertian malaria** (fever spikes 48h) :
 - 1. Immature trophozoite/ **ring** (nuclear material resembles diamond)
 - 2a. **Mature trophozoite**
 - 2b. Development into gametocytes - sexual reproduction if ingested by anopheles;
 - 3. **Schizont**, rupture RBC releasing **merozoites** -> **fever, chills and sweats**

Plasmodium falciparum

- Tropical and subtropical regions with **frequent co-infection with HIV**
 - **shortest incubation period**: 7-10 days
 - early influenza-like signs
 - chills, fever with profuse sweating
 - **malignant tertian malaria** (fever spikes 36-48h)
 - capillary plugging
 - **cerebral malaria that can lead to coma and death**
 - **kidney damage** – intravascular hemolysis and hemoglobinuria
 - **hepatosplenomegaly** – abdominal pain, vomiting of bile, diarrhea and dehydration
- trophozoite/ schizonts in liver & spleen**
- **peripheral blood smears: only young rings/ gametocytes**

Plasmodium falciparum

- Chloroquine-resistant strains in all endemic areas (Africa, Southeast Asia, South America); exception in Central America & Caribbean regions
- If not-resistant: **chloroquine** or **parenteral quinine**;
- **Resistant:**
 - mefloquine ± artesunate
 - artemether ± lumefantrine
 - quinine, quinidine, doxycycline, atovaquone-proguani, pyrimethamine-sulfadoxine
quinine, pyrimethamine-sulfadoxine: may be toxic; treatment rather than prophylaxis
- **Newer agents** against multidrug-resistant strains:
 1. Phenanthrene methanols: halofantrine and lumefantrine;
 2. Artemisinins: artemether, artesunate
- * **Whole-blood exchange: if available**; in severe cases; cerebral malaria, acute lung injury, severe hemolysis with acidemia, shock, or a rising level of parasitemia despite adequate IV therapy

P. vivax and *P. ovale*


- Differences from *P. falciparum*:
- Most prevalent; in tropics, subtropics and temperate regions
- Incubation 10-17 days
- **Tertian malaria** – chills, fever, sweats every 48h
- Ability to form **hypnozoites** in liver – **cause relapsing malaria!**
- **Treatment:** Chloroquine + Primaquine (for hypnozoites)

P. malariae


- Longest Incubation: 18-40days
- **Quartan malaria – 72h intervals**
- **Moderate to severe attacks** and can persist for hours

Plasmodium - diagnosis


- **Thick drop and thin blood smears** (e.g. fig. 1- *P. falciparum* trophozoites as headphones in thick drop and fig. 2-blood smear) 1000x under oil immersion;
- infected RBCs do not enlarge/ become distorted (vs *P. vivax* and *P. ovale*)
- **PCR diagnosis from venous blood** (fig. 3)
- **Rapid Ag based diagnostic tests** (fig. 4)



https://www.researchgate.net/figure/Giemsa-stained-thick-film-with-high-parasite-density-of-P-falciparum-Photograph-by_fig2_227180663



<https://www.cellwiki.net/en/aberrations/microorganisms-malaria-falciparum>



<https://www.globalpointofcare.abbott/ww/en/product-details/binaxnow-malaria.html>

Natural protection

- Absence of RBC membrane antigens **Duffy** that *P. vivax* uses for binding
- Sickle cell anemia trait protects RBC from *P. falciparum*